

QFT and its discontents
a blog

Predrag Cvitanović

August 20, 2018

Contents

| | | |
|----------|---|-----------|
| 1 | Is QED finite? | 3 |
| 1.1 | Electron magnetic moment | 3 |
| 1.2 | Gauge sets | 5 |
| 1.2.1 | QED vertex photons come in three “colors” | 7 |
| 1.2.2 | The unreasonable smallness of gauge sets | 8 |
| 1.2.3 | Self-energy sets | 9 |
| 1.2.4 | Where do we go from here? | 9 |
| 1.3 | Worldline formalism | 13 |
| 1.3.1 | High-orders QED in worldline formalism | 16 |
| 1.3.2 | Electron magnetic moment in worldline formalism | 17 |
| 1.4 | Volkov method | 19 |
| 1.5 | Hopf algebraic approach | 20 |
| 1.6 | Method of smooth conjugacies | 20 |
| 1.7 | Summary | 24 |
| 1.8 | Is QED finite? Correspondence | 25 |
| 1.9 | QCD gauge sets - a blog | 37 |
| 1.10 | Is QED finite? A blog | 42 |
| 1.10.1 | Quenched QED/QFT | 42 |
| 1.10.2 | Is QED finite? A blog, continued | 43 |
| | References | 61 |
| 2 | Learning worldline QFT | 72 |
| 2.1 | Quantum Field Theory Spring 2018 Report | 72 |
| 2.2 | Morelia Worldline Formalism course | 72 |
| 2.3 | Xu wordline formalism notes | 76 |
| | References | 82 |
| 3 | Daily QFT blog | 84 |
| | References | 99 |

Chapter 1

Is QED finite?

For Emily, after she gets bored with “Baby Loves Quarks.”

In 2017 Laporta [116, 117] completed the twenty-year project of computing analytically the individual contributions of 891 4-loop vertex diagrams contributing to the electron magnetic moment g . Vertex diagrams separate in 25 gauge-invariant sets (see figure 1.8). The numerical contribution of each set is listed in table 1.2. Adding only the quenched set V diagrams (diagrams with no lepton loops, see figure 1.4 and 1.6), one finds for the 4- and 5-loop contributions to the anomaly $a[V] = \frac{1}{2}(g - 2)|_V$:

$$\begin{aligned} a^{(8)}[V] &= -2.176866027739540077443259355895893938670 \\ &= -2.17 \dots && \text{Aoyama } et al. \text{ 2012 [7]} \\ &\approx 0 && \text{Cvitanović 1977 [41]} \\ a^{(10)}[V] &= 7.606(192) \dots && \text{Aoyama } et al. \text{ 2018 [11]} \\ &\approx 3/2 && \text{Cvitanović 1977 [41]}. \end{aligned} \tag{1.1}$$

There is a prediction dating back to 1977 for values of these terms: the predicted $a^{(8)}[V] \approx 0$ does not pan out, but the difference is small, considering that this is a sum of 518 vertex diagrams (or 47 self-energy diagrams) [105]. Likewise, the prediction for $a^{(10)}[V]$ is not too far off, considering that this is a sum of 6 354 vertex diagrams of table 1.1 (or 389 self-energy diagrams).

1.1 Electron magnetic moment

This section sets up the notation - the reader can safely skip it and start with sect. 1.2. For an introduction to the conventional magnetic moment calculation, see for example the Cvitanović online graduate QFT course, lectures 25 and 26 [here](#).

An electron of mass m has a magnetic moment

$$\mu = \frac{e\hbar}{2mc} \frac{g}{2} \tag{1.2}$$

where g is the gyromagnetic ratio. In Dirac theory [54], the electron has $g = 2$.

Consider the electron-photon vertex Γ_μ of quantum electrodynamics, with $p_i = p - q/2$ and $p_o = p + q/2$ the momenta of incoming and outgoing electron lines, evaluated on the electron mass shell $p_i^2 = p_o^2 = m^2$. By Gordon decomposition the vertex can be written in terms of the Dirac and Pauli form factors $F_1(q^2)$ and $F_2(q^2)$:

$$\bar{u}(p_o)\Gamma_\mu(p, q)u(p_i) = \bar{u}(p_o)\left\{F_1(q^2)\gamma_\mu - \frac{F_2(q^2)}{2m}\sigma_{\mu\nu}q^\nu\right\}u(p_i), \quad (1.3)$$

where the spinors $\bar{u}(p_i)$ and $u(p_i)$ satisfy the Dirac equation:

$$\bar{u}(p_o)\not{p}_o = m\bar{u}(p_o), \quad \not{p}_i u(p_i) = m u(p_i).$$

We follow the notation of Bjorken and Drell [21] and Cvitanović and Kinoshita [50]. Z_1 , Z_2 , and Z_3 , are the respectively the vertex, the electron wave function, and the photon wave function renormalization constants, and the electron mass will be set to $m = 1$ throughout. In what follows it is convenient to define $Z_1 = 1 + L$. For QED the charge conservation requires that the renormalized charge form factor satisfies $\tilde{F}_1(0) = 1$, which is guaranteed by the Ward identity [180] $Z_1 = Z_2$. The vertex renormalization constant L is given by the on-shell value of the unrenormalized charge form factor [30]

$$1 + L = F_1(0) = \frac{1}{4}\text{tr}[(\not{p} + 1)p^\nu\Gamma_\nu]_{q=0}, \quad (1.4)$$

and $a = (g - 2)/2$, the anomalous magnetic moment of an electron is given by the static limit of the magnetic form factor $a = \tilde{F}_2(0) = M/(1 + L)$, where [30]

$$M = \lim_{q \rightarrow 0} \frac{1}{4q^2} \text{tr} \left\{ [\gamma^\nu p^2 - (1 + q^2/2)p^\nu] (\not{p}_o + 1)\Gamma_\nu(\not{p}_i + 1) \right\}. \quad (1.5)$$

The perturbative expansions for the magnetic moment anomaly is defined as

$$a = \frac{M(\alpha_0)}{1 + L(\alpha_0)} = \sum_{n=1}^{\infty} a_0^{(2n)} \left(\frac{\alpha_0}{\pi}\right)^n, \quad (1.6)$$

where $1 + L = F_1(0)$, $M = F_2(0)$ are computed from the unrenormalized proper vertex (1.3), given by the sum of all one-particle irreducible electron-electron-photon vertex diagrams with internal photons, electron loops and electron mass counterterms. Expanding M and L we have

$$\begin{aligned} a_0^{(2)} &= M^{(2)} \\ a_0^{(4)} &= M^{(4)} - L^{(2)}M^{(2)} \\ a_0^{(6)} &= M^{(6)} - L^{(2)}M^{(4)} - (L^{(4)} - (L^{(2)})^2)M^{(2)} \end{aligned} \quad (1.7)$$

As shown in ref. [39], for the anomaly (1.6) expressed in terms of the unrenormalized coupling constant α_0 , all $a_0^{(n)}$ are IR finite, for both QED and QCD. The UV finite expression for the anomaly (1.6) is obtained by the charge renormalization

$$\alpha = Z\alpha_0, \quad Z = \frac{Z_2}{Z_1} Z_3, \quad (1.8)$$

| Order $2n$ | Vertex graphs Γ_{2n} | Gauge sets G_{2n} | Anomaly $a^{(2n)}$ |
|---------------|--------------------------------|------------------------|-----------------------|
| 2 | 1 | 1 | 1/2 |
| 4 | 6 | 2 | 0 |
| 6 | 50 | 4 | 1 |
| 8 | 518 | 6 | 0 |
| 10 | 6354 | 9 | 3/2 |
| 12 | 89 782 | 12 | 0 |
| 14 | 1 429 480 | 16 | 2 |

Table 1.1: Comparison of the number of vertex diagrams without fermion loops, gauge sets, and the gauge-set approximation (1.12) for the magnetic moment in $2n$ th order. From ref. [41].

where Z_1 , Z_2 , and Z_3 are computed as power series in the bare coupling constant α_0 . For QED $Z_1 = Z_2$ by the Ward identity [180], and for QCD the Z_i 's are related by the Taylor-Slavnov [164, 173] identities (1.8). The simple structure of (1.6) and (1.7) should simplify the worldline calculations of sect. 1.3.2.

The Dirac equation predicts that the magnetic moment of an electron of charge e and mass m is $\mu = e/2m$, i.e., in the absence of radiative corrections $\tilde{F}_1(0) = 1$ and $\tilde{F}_2(0) = 0$. In 1948 Schwinger [156] showed that in the leading, one-loop order in the fine structure constant α , the radiative corrections lead to the anomalous magnetic moment of form $\tilde{F}_2(0) = \alpha/2\pi + a^{(2)} (\alpha/\pi)^2 + \dots$ (the result engraved on Schwinger's tombstone). These notes are about what to expect for the $(\alpha/\pi)^n$ term in this series.

1.2 Gauge sets

Is there any method of computing the anomalous moment of the electron which, on first approximation, gives a fair approximation to the α term and a crude one to α^2 ; and when improved, increases the accuracy of the α^2 term, yielding a rough estimate to α^3 and beyond?

— Feynman's challenge, 12th Solvay Conference [70]

In 1972 Toichiro Kinoshita and Predrag Cvitanović had completed computing a large number of 3-loop anomalous magnetic moment Feynman diagrams and regularization counterterms [104], figure 1.1. The subsequent 4- and 5-loop numerical and analytic calculations were nothing short of heroic [9, 105, 116, 118]. The quantum field theory was used in the standard way [21], by expanding the magnetic moment into combinatorially many Feynman diagrams (see the numbers of vertex graphs in table 1.1). Each Feynman diagram corresponds to an integral in many dimensions, with oscillatory integrand with thousands of terms, each integral separately UV divergent, IR divergent,




Figure 1.1: The three-loop vertex diagrams contributing to $A_1^{(6)}$ magnetic moment (from Jegerlehner and Nyffeler [92]). Lautrup *et al.* [122] were the first to note that subsets $(3, 0, 0) = \{23, 24, 25, 26, 27, 28\}$; $(2, 1, 0) = \{29, 31, 33, 35, 37, 39, 41, 43, 45, 47\}$ and its time-reversal $(2, 0, 1) = \{30, 32, 34, 36, 38, 40, 42, 44, 46, 48\}$; $(1, 2, 0) = \{49, 51, 53, 55, 57, 59, 61, 63, 65, 67\}$ and its time-reversal $(1, 0, 2) = \{50, 52, 54, 56, 58, 60, 62, 64, 66, 68\}$; and $(1, 1, 1) = \{69, 70, 71, 72\}$ are the minimal gauge sets, see figure 1.3.




Figure 1.2: Rows: the fourth-order gauge sets $km'm$: (1) = (1, 1, 0), (2) = (2, 0, 0) and (3) = (1, 0, 1). Columns: external field insertions into the two self-energy sets. For diagrams related by time reversal (here (1) and (3)) the value listed under the first diagram of the pair is the total contribution of the pair. Contributions seem to be of order $\pm \frac{1}{3} \left(\frac{\alpha}{\pi}\right)^2$, and suggest that a set and its time-reversed partner should be counted separately. From ref. [41].

and unphysical, as its value depends on the definition of counterterms and the choice of gauge. The numerical values of these integrals typically range from ± 10 to ± 100 . For example, the largest contributions of the 389 quenched self-energy diagrams listed in Aoyama *et al.* 2018 [11] are of order ± 20 .

Adding up hundreds of such contributions, of wildly fluctuating values, yields (for the no-fermion loops subset V , in the notation of ref. [9])

$$A_1^{(6)}[V] = +(0.92 \pm 0.02) \left(\frac{\alpha}{\pi}\right)^3 .$$

But why “+” and not “-”? Why so small? Why does a sum of hundreds of diagrams and counterterms yield a number of order of unity, and not 10 or 100 or any other number?

If gauge invariance of QED guarantees that all UV and on-mass shell IR divergences cancel, could it be that it also enforces cancellations among the finite parts of contributions of different Feynman graphs?

1.2.1 QED vertex photons come in three “colors”

As first noted by Lautrup, Peterman and de Rafael [122], the renormalized on-mass shell QED vertex diagrams separate into a sum of minimal gauge-invariant subsets, each subset separately UV and IR finite. The only published proof of this elementary fact seems to be ref. [41]. The very reasonably priced ref. [43] might be worth a

read, especially if one is interested in the non-abelian theories as well [39, 40, 42], see sect. 1.9.

A gauge change generates a k^μ term in a photon propagator, and that affects a tree electron vertex in a very simple way, $\not{k} = (\not{p} + \not{k} + m) - (\not{p} + m)$, or, diagrammatically [21, 43] (graphs drawn by H. Kibler [107]),¹

$$\frac{1}{\not{p} + \not{k} - m} \not{k} \frac{1}{\not{p} - m} = \frac{1}{\not{p} - m} - \frac{1}{\not{p} + \not{k} - m}, \quad (1.9)$$

To simplify matters, in what follows we shall consider only the no-fermion loop diagrams, or ‘quenched-’, or ‘q-type’ diagrams (‘quenched’, as this corresponds to the N_f -independent part of the vertex amplitude in QED with N_f flavors). The minimal gauge-invariant subsets without electron loops (see figure 1.1 diagrams {23 – 72}; figure 1.2; 1.3; 1.4; and 1.6) will be hereafter be referred to as *gauge sets*.

A gauge set $km'm$ consists of all 1-particle irreducible vertex diagrams without electron loops, with k photons crossing the external vertex (cross-photons) and $m[m']$ photons originating and terminating on the incoming [outgoing] electron leg (leg-photons), where $m \geq m'$. For asymmetric pairs of sets, with $m \neq m'$, the contribution to the anomaly $a_{kmm'}$ is, in the convention of ref. [41], the sum of the set and its mirror (time-reversed) image,

$$a[V] = \frac{1}{2}(g-2) \Big|_V = \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} \sum_{m'=0}^m a_{kmm'} \left(\frac{\alpha}{\pi}\right)^{k+m+m'}. \quad (1.10)$$

1.2.2 The unreasonable smallness of gauge sets

When the diagrams computed in ref. [50] are grouped into gauge sets, figure 1.2 to figure 1.6, a surprising thing happens; while the finite part of each Feynman diagram is of order of 10 to 100, every gauge set known at the time added up to approximately

$$\pm \frac{1}{2} \left(\frac{\alpha}{\pi}\right)^n,$$

with the sign given by a simple empirical rule

$$a_{kmm'} = (-1)^{m+m'} \frac{1}{2}. \quad (1.11)$$

The sign rule is further corroborated by sets with photon self-energy insertions (but with the absolute size scaled down to 3 – 15% of (1.11)). In figure 1.4 this rule is compared with the actual numbers, and the 1977 four-loop prediction is given [41]. With that prediction, the “zeroth” order estimate of the electron magnetic moment anomaly a is given by the “gauge-set approximation,” convergent and summable to all orders

$$a = \frac{1}{2}(g-2) = \frac{1}{2} \frac{\alpha}{\pi} \frac{1}{\left(1 - \left(\frac{\alpha}{\pi}\right)^2\right)^2} + \text{“corrections”}. \quad (1.12)$$

¹Predrag: 2017-07-14 need to download, put axohelp.exe, the executable version of axohelp for MS-Windows into the directory where I can run it from, presumably Program Files/MiKTeX 2.9/miktex/bin/x64

This is not how one usually thinks of perturbation theory. Most of our colleagues believe that in 1952 Dyson [61] had shown that the perturbation expansion is an asymptotic series (for a discussion, see Dunne and Schubert [60, 89]), in the sense that the n -th order contribution should be exploding combinatorially

$$\frac{1}{2}(g-2) \approx \dots + n^n \left(\frac{\alpha}{\pi}\right)^n + \dots,$$

and not growing slowly like my estimate

$$\frac{1}{2}(g-2) \approx \dots + \frac{n}{2} \left(\frac{\alpha}{\pi}\right)^{2n} + \dots.$$

For me, the above is the most intriguing hint that something deeper than what we know today underlies quantum field theory, and the most suggestive lesson of our calculation.

1.2.3 Self-energy sets

There are two ways of grouping vertex diagrams, into *gauge sets*, and into *self-energy sets* (or the “externally gauge-invariant” sets). Every vertex diagram belongs both to a gauge set and to a self-energy set, as illustrated by figure 1.3. Formulation of the $(g-2)$ computation directly from self-energy graphs is due to Cvitanović and Kinoshita, see the “new formula” (6.22) in ref. [50]. Not only does this calculation use fewer Feynman graphs, but it enables calculation of the 3-loop electron magnetic moment by two independent methods (and in this way helps track down and eliminate errors in either calculation). As an important aside, Carroll [31, 32] gives the credit for self-energy sets only to the mass-operator formalism of Schwinger [157–159, 166], even though Carroll papers look closer to ref. [50] than to Schwinger and Sommerfield; ref. [50] is cited, and his derivation is also based on the Ward-Takahashi identity [172, 180]. The self-energy set formulation might be equivalent to Schwinger’s, but it looks quite different in detail, and the authors were not aware of Schwinger mass-operator when they derived it.

The gauge sets are minimal, and separately gauge invariant (for a proof, see ref. [41]). The self-energy sets are not, only their sum is gauge invariant. Unlike gauge sets, whose number grows polynomially, the number of self-energy sets grows combinatorially - they save significant amount of computing for few-loops computations, but cannot be used to argue the finiteness of QED. That is the reason why Aoyama *et al.* [7, 9, 11] calculations have nothing to say about the 1977 conjecture [41]: they do not compute individual vertex diagrams, but only the self-energy sets, and for them the set of all diagrams without a fermion loop (‘quenched-’ or ‘q-type’ diagrams) is a single ‘gauge-invariant set’ V . For example, for 5-loops the set V is a sum of 9 vertex gauge sets (where time-reversed pairs count as one set, see figure 1.5), but Aoyama *et al.* [11] only give their sum (1.1).

1.2.4 Where do we go from here?

Gauge invariance is the bane of my life

— Predrag




Figure 1.3: Every vertex diagram belongs both to a 'gauge set' and to a 'self-energy set'. This table illustrates the two kinds of sets. The 3-loop gauge sets $kim'm$ are arranged in the rows, and the self-energy sets (or the 'externally gauge-invariant' sets, vertex diagrams obtained by inserting an extra vertex into a self-energy diagram) in the columns, labeled as in Fig. 3 of ref. [50]. The values are finite parts in the $\ln \lambda$ IR cut-off approach, such as those listed in ref. [124]. For different IR separation methods (such as in ref. [50]) and different gauges, individual diagrams have different values. The gauge sets, however, are separately gauge invariant. The self-energy sets (whose number grows combinatorially with the order in perturbation theory) are not, only their sum is gauge invariant. From ref. [41].



| $2n$ | | anomaly |
|------|---|---------|
| 2 |  $\frac{1}{2}$ | 5 |
| 4 |  $-\frac{1}{2} (-65)$  $\frac{1}{2} (31)$ | 0 (-33) |
| 6 |  $\frac{1}{2} (56)$  $-\frac{1}{2} (-47)$  $\frac{1}{2} (43)$ | 1 (93) |
| |  $\frac{1}{2} (44)$ | |
| 8 |  $-\frac{1}{2}$  $\frac{1}{2}$  $-\frac{1}{2}$  $\frac{1}{2}$ | 0 (?) |
| |  $-\frac{1}{2}$  $\frac{1}{2}$ | |

Figure 1.4: Comparison of the 1977 gauge-set approximation to the anomaly a and the actual numerical values of corresponding gauge sets, together with the 1977 eighth-order prediction of ref. [41]. For the updated listing, see figure 1.5.

| $2n$ | $km'm$ | | | | | anomaly |
|------|--|---|--|---|---|----------------------|
| 2 | $(\mathbf{1}, \mathbf{0}, \mathbf{0})$ $1/2$ | | | | | $\frac{1}{2}$ |
| 4 | $(\mathbf{1}, \mathbf{1}, \mathbf{0})$ $-1/2 (-.65)$ | $(\mathbf{2}, \mathbf{0}, \mathbf{0})$ $1/2 (.31)$ | | | | 0 (-.33) |
| 6 | $(\mathbf{1}, \mathbf{2}, \mathbf{0})$ $1/2 (.56)$ | $(\mathbf{2}, \mathbf{1}, \mathbf{0})$ $-1/2 (-.47)$ | $(\mathbf{3}, \mathbf{0}, \mathbf{0})$ $1/2 (.44)$ | | | 1 (.93) |
| 8 | $(\mathbf{1}, \mathbf{1}, \mathbf{1})$ $1/2 (.43)$ | | | | | |
| 8 | $(\mathbf{1}, \mathbf{3}, \mathbf{0})$ $-1/2 (-1.97)$ | $(\mathbf{2}, \mathbf{2}, \mathbf{0})$ $1/2 (-.14)$ | $(\mathbf{3}, \mathbf{1}, \mathbf{0})$ $-1/2 (-1.04)$ | $(\mathbf{4}, \mathbf{0}, \mathbf{0})$ $1/2 (.51)$ | 0 (-2.17) | |
| 10 | $(\mathbf{1}, \mathbf{2}, \mathbf{1})$ $-1/2 (-.62)$ | $(\mathbf{2}, \mathbf{1}, \mathbf{1})$ $1/2 (1.08)$ | | | | |
| 10 | $(\mathbf{1}, \mathbf{4}, \mathbf{0})$ $1/2 (?)$ | $(\mathbf{2}, \mathbf{3}, \mathbf{0})$ $-1/2 (?)$ | $(\mathbf{3}, \mathbf{2}, \mathbf{0})$ $1/2 (?)$ | $(\mathbf{4}, \mathbf{1}, \mathbf{0})$ $-1/2 (?)$ | $(\mathbf{5}, \mathbf{0}, \mathbf{0})$ $1/2 (?)$ | $\frac{3}{2}$ (7.60) |
| 10 | $(\mathbf{1}, \mathbf{3}, \mathbf{1})$ $1/2 (?)$ | $(\mathbf{2}, \mathbf{2}, \mathbf{1})$ $-1/2 (?)$ | $(\mathbf{3}, \mathbf{1}, \mathbf{1})$ $1/2 (?)$ | | | |
| 10 | $(\mathbf{1}, \mathbf{2}, \mathbf{2})$ $1/2 (?)$ | | | | | |

Figure 1.5: Updated figure 1.4 comparison of the gauge-set approximation (1.12) and the actual numerical values of corresponding gauge sets, together with the 5-loop prediction. Starting with 4-loops, the gauge-set approximation fails in detail. Still, the signs are right, except for the anomalously small set $(2, 2, 0)$, and the remaining sets are surprisingly close to multiples of $1/2$.



| gauge set | $km'm$ | value | prediction |
|-----------|---------|----------|------------|
| (1) | (1,3,0) | - 1.9710 | - 1/2 |
| (2) | (2,2,0) | - 0.1424 | 1/2 (!) |
| (3) | (1,2,1) | - 0.6219 | - 1/2 |
| (4) | (2,1,1) | 1.0867 | 1/2 |
| (5) | (3,1,0) | - 1.0405 | - 1/2 |
| (6) | (4,0,0) | 0.5125 | 1/2 |

Figure 1.6: (top) Examples of 4-loop vertex diagrams belonging to Laporta [116] gauge sets (1) to (6). The remaining diagrams in the set can be obtained by permuting separately the vertices on the left and right side of the electron line, and considering also the mirror images of the diagrams. For all 25 gauge-invariant sets, see figure 1.8. The table: Gauge-set contributions $a_{kmm'}^{(8)}$, see (1.10), as reported by Laporta [116] (for the full 25 gauge-invariant sets, see table 1.2). The last column: 1977 Cvitanović predictions [41]. Signs are right, except for the set (2) = (2, 2, 0), which is anomalously small, and the remaining sets are surprisingly close to multiples of 1/2. There might be factors of 2 having to do with symmetries, missing from the guesses of ref. [41], but I cannot see how that would work. Only (4) = (2, 1, 1) and (6) = (4, 0, 0) are symmetric, but (1) = (1, 3, 0), (4) and (5) = (3, 1, 0) seem to have an extra factor of 2 or 4.

Aoyama *et al.* 5-loop calculations already push the envelope of what is numerically attainable, they cannot switch from the self-energy diagrams formulation to the vertex diagrams formulation, it would mean (for the quenched set V) going from 389 self-energy graphs to 6354 vertex diagrams. Stefano Laporta deserves a bit of well earned rest. So what is ahead?

At this time, Sergey A. Volkov appears to be the only person set up do the requisite 5-loop calculations, see sect. 1.4.

Two approaches might be relevant to establishing bounds on, and perhaps even the direct computation of gauge sets (ignoring the $N = 2$ and $N = 4$ supersymmetric models): (1) *worldline formalism* pursued by Schubert and collaborators, see sect. 1.3, and (2) *Hopf algebraic approach* of Kreimer and collaborators, see sect. 1.5.

Very far out in the left field is the smooth conjugacy method of sect. 1.6 which would require a bit of real work to apply it to a field theory.

1.3 Worldline formalism

How and why Feynman in 1950 introduced ‘worldline formalism’ (initially for scalar QED, appendix to ref. [67], then for spinor QED, appendix to ref. [68]) is explained

in Schubert 2001 report [152] (which also has an extensive bibliography up to 2001). In 1982 Affleck, Alvarez, and Manton [1] used the Feynman worldline path integral representation of the quenched effective action for scalar QED in the constant electric field.

For the remainder of this section we shall consider only the quenched QED, i.e., restrict our considerations only to sets of diagrams with no lepton loop insertions.

A formula for the charged scalar propagator to emit and reabsorb N photons as it propagates from x' to x can be derived as follows [3]. The free scalar propagator for the Euclidean Klein-Gordon equation [3, 153] is

$$D_0(x, x') = \langle x | \frac{1}{-\square + m^2} | x' \rangle, \quad (1.13)$$

where \square is the D -dimensional Laplacian. Exponentiate the denominator following Schwinger,

$$D_0(x, x') = \int_0^\infty dT e^{-m^2 T} \langle x | e^{-T(-\square)} | x' \rangle, \quad (1.14)$$

Replace the operator in the exponent by a path integral

$$D_0(x, x') = \int_0^\infty dT e^{-m^2 T} \int_{x(0)=x'}^{x(T)=x} \mathcal{D}x(\tau) e^{-\int_0^T d\tau \frac{1}{4} \dot{x}^2}, \quad (1.15)$$

where τ is a proper-time parameter (the fifth parameter [71]), and the dot denotes a derivative with respect to the proper time. This is the *worldline path integral* representation of the relativistic propagator of a scalar particle in Euclidean space-time. In the vacuum (no background field), it is easily evaluated by standard methods and leads to the usual space and momentum space free propagators,²

$$\int_{x(0)=x'}^{x(T)=x} \mathcal{D}x(\tau) e^{-\int_0^T d\tau \frac{1}{4} \dot{x}^2} = \frac{1}{(4\pi T)^{d/2}}. \quad (1.16)$$

Adding the QED interaction terms leads to the Feynman's worldline path integral representation [67] of the charged scalar propagator of mass m in the presence of a background field $A(x)$,

$$D(x, x') = \int_0^\infty dT e^{-m^2 T} \int_{x(0)=x'}^{x(T)=x} \mathcal{D}x(\tau) e^{-S_0 - S_e - S_i}, \quad (1.17)$$

where the suffix (0) indicates the free propagation

$$S_0 = \int_0^T d\tau \frac{1}{4} \dot{x}^2, \quad (1.18)$$

(e) is the interaction of the charged scalar with the external field

$$S_e = -ie \int_0^T d\tau \dot{x}^\mu A_\mu(x(\tau)), \quad (1.19)$$

²Predrag: 2017-06-17 Here a study of Sect. 6. *Worldline formalism* of Gelis and N. Tanji [80] might be helpful - it reexpresses the integral as an average over Wilson loops.

and (i) are the virtual photons exchanged along the charged particle's trajectory

$$S_i = \frac{e^2}{2} \int_0^T d\tau_1 \int_0^T d\tau_2 \dot{x}_1^\mu D_{\mu\nu}(x_1 - x_2) \dot{x}_2^\nu, \quad (1.20)$$

where $D_{\mu\nu}$ is the x -space photon propagator.

The formula (1.17) involves (i) a path integral over all the worldlines $x(\tau')$, i.e., closed paths in Euclidean space-time parameterized by the proper time $\tau' \in [0, \tau]$, and (ii) an ordinary integral over the length τ of these paths. The sum over all the worldlines accounts for the quantum fluctuations in space-time, and the prefactor $\exp(-m^2 T)$ suppresses the very long worldlines that explore regions of space-time much larger than the Compton wavelength of the particles. The ultraviolet properties of the theory are encoded in the short worldlines limit $\tau \rightarrow 0$. The Euclidean space-time guarantees that both types of integrals are convergent.

Consider next the charged scalar field in external field, neglecting internal photon loops. By taking the constant external field $A(x)$ to be a sum of N plane waves, one obtains the rule for inserting N external photons:

$$\begin{aligned} D_{(N)}(x, x') &= (-\lambda)^N \int_0^\infty dT e^{-m^2 T} \int_0^T d\tau_1 \cdots \int_0^T d\tau_N \\ &\times \int_{x(0)=y}^{x(T)=x} \mathcal{D}x e^{i \sum_{i=1}^N k_i \cdot x(\tau_i)} e^{-\int_0^T d\tau \frac{1}{4} \dot{x}^2}. \end{aligned} \quad (1.21)$$

For the spinor case, the magnetic moment will be given by the term linear in a constant external field $A(x)$, and in order to define gauge sets, one will have to distinguish the in- and out-electron lines.

The object of great interest to us is the quenched internal virtual photons term (1.20):

$$\int_{x(0)=x'}^{x(T)=x} \mathcal{D}x(\tau) e^{-S_i} = \int_{x(0)=x'}^{x(T)=x} \mathcal{D}x(\tau) e^{-\frac{e^2}{2} \int_0^T d\tau_1 \int_0^T d\tau_2 \dot{x}_1^\mu D_{\mu\nu}(x_1 - x_2) \dot{x}_2^\nu}. \quad (1.22)$$

(Fried and Gabellini [77] refer to this as the ‘‘linkage operator’’). Expanded perturbatively in α/π , this yields the usual Feynman-parametric vertex diagrams. However, it is Gaussian in \dot{x}^μ , and if by integration by parts, \dot{x}^μ are eliminated in favor of x^μ , internal photons can be integrated over directly, prior to an expansion in $(\alpha/\pi)^n$, and one gets integrals in terms of N -photon propagators, symmetrized sums over N photons, and not the usual Feynman graphs. Each usual Feynman graph corresponds to one particular permutation of internal photon insertions, and from that comes the factorial growth in the number of graphs.

These integrations by parts lead to the first and second proper-time derivatives of the Green's function, worked out in the literature (for example, in refs. [153, 169]), the details would take too much space to recap here. I find Bastianelli, Huet, Schubert, Thakur and Weber 2014 paper [17] quite inspirational. My notes on these papers are below, around page 45. Apologies, my notes are just a jumble, jottings taken as I try to understand this literature. They might be useful anyway, as pointers to the literature.




Figure 1.7: Quenched diagrams contributing to the three loop QED photon propagator. From ref. [17].

Thus, for the quenched scalar QED, the worldline integrals are expressed in terms of N -photon propagators, the central ingredient that defines the quenched gauge sets (1.10). Unlike the Feynman parameter integrals for individual vertex graphs, they are independent of the ordering of the momenta k_1, \dots, k_N ; the formula (1.22) contains all $\approx N!$ ways of attaching the N photons to the charged particle propagator. The formulation combines combinatorially many Feynman diagrams into a single integral. An example are the quenched contributions to the three-loop photon propagator shown in figure 1.7.

In QED the N -photon propagator formulation combines into one integral all Feynman graphs related by permutations of photon legs along fermion lines, that is, it should yield *one* integral for a gauge set $km'm$ defined in (1.10).

1.3.1 High-orders QED in worldline formalism

A non-perturbative formula for QED in a constant field, given for scalar QED in 1982 by Affleck, Alvarez, and Manton [1] is an example how the worldline formalism can yield high-order information on QED amplitudes. Huet, McKeon, and Schubert [88] continue this in their 2010 study of the 1-electron loop, N -photon amplitudes in the limit of large photon numbers and low photon energies, this time for 1+1 dimensional scalar QED, in order to illustrate the large cancellations inside gauge invariant classes of graphs.

Affleck *et al.* [1] use the Feynman [67] ‘worldline path integral’ representation of the quenched effective action for scalar QED in the constant electric field, and calculate the amplitude in a stationary path approximation. The stationary trajectory so obtained is a circle with a field dependent radius, called “instanton” in this context. The worldline action on this trajectory yields the correct exponent, and the second variation determinant yields the correct prefactor. Using Borel analysis, they obtain non-perturbative information on the on-shell renormalized N -photon amplitudes at large N and low energies.

Parenthetically, independently and not by worldline formalism, but by dint of difficult calculations and much deep physics intuition, Lebedev and Ritus [123] have arrived at a nonperturbative mass shift interpretation for the spinor QED pair creation in constant electric field in 1984.

For the quenched spinor QED (fermion lines decorated by photon exchanges) closed-form expressions for general N require the worldline super-formalism [152], at the cost of introducing Fradkin 1966 [72] Grassmann path integral, or, alternatively, the

second order formalism of Strassler [169].

G. Torgrimsson, Schneider, Oertel and Schützhold [175] 2017, [arXiv:1703.09203](#), sect. 3: use the N -photon formalism to determine a saddle and the asymptotic form of two types of dynamically assisted Sauter-Schwinger effect.

The 2006 Dunne and Schubert [60] study of scalar and spinor QED N -photon amplitudes, in the quenched approximation (i.e., taking only the diagrams with one electron loop) led to “the following generalization of Cvitanović’s conjecture: the perturbation series converges for all on-shell renormalized QED amplitudes at leading order in N_f . It must be emphasized that the on-shell renormalization is essential in all of the above.” Unlike Cvitanović [41] purely numerical conjecture, theirs is a sophisticated argument, buttressed by Borel dispersion relations.

1.3.2 Electron magnetic moment in worldline formalism

Here we specialize the electron magnetic moment discussion of sect. 1.1 to the quenched subsector. Z_1 , Z_2 , and Z_3 , are the respectively the vertex, the electron wave function, and the photon wave function renormalization constants. For quenched QED there are no fermion loops, there are no vacuum polarization contribution to the charge renormalization (1.8), $Z = Z_3 = 1$, so the bare coupling equals the physical coupling, $\alpha_0 = \alpha$. Furthermore, $Z_1 = Z_2$ by the Ward identity [180].

The anomalous magnetic moment of an electron $a = (g - 2)/2$ is given by the static limit of the magnetic form factor $a = \tilde{F}_2(0) = M/(1 + L)$ from (1.5), with perturbative expansion

$$a = \frac{M(\alpha)}{1 + L(\alpha)} = \sum_{n=1}^{\infty} a^{(2n)} \left(\frac{\alpha}{\pi}\right)^n, \quad (1.23)$$

where $Z_1 = 1 + L = F_1(0)$, $M = F_2(0)$ are computed from the unrenormalized on-shell values of proper vertex (1.3), given by the sum of all one-particle irreducible (1pI) electron-electron-photon vertex diagrams with internal photon corrections (no electron loops). Expanding M and L we have

$$\begin{aligned} a_0^{(2)} &= M^{(2)} \\ a_0^{(4)} &= M^{(4)} - L^{(2)}M^{(2)} \\ a_0^{(6)} &= M^{(6)} - L^{(2)}M^{(4)} - (L^{(4)} - (L^{(2)})^2)M^{(2)} \end{aligned} \quad (1.24)$$

Each order in (1.23) is IR and UV finite, with the UV subdivergences are cancelled by $L^{(2m)}$ counterterms in (1.24).

A gauge set $km'm$ in expansion (1.10) consists of all 1-particle irreducible vertex diagrams without electron loops, with k photons crossing the external vertex (cross-photons) and $m[m']$ photons originating and terminating on the incoming [outgoing] electron leg (leg-photons). One can assume three different coupling, setting them all equal to α at the end of the calculation,

$$a = \sum_{m'=0}^{\infty} \left(\frac{\alpha'}{\pi}\right)^{m'} \sum_{k=1}^{\infty} \left(\frac{\alpha_v}{\pi}\right)^k \sum_{m=0}^{\infty} \left(\frac{\alpha}{\pi}\right)^m a_{km'm}. \quad (1.25)$$

The gauge set contributions are then

$$\begin{aligned}
a^{(2)} &= a_{100} \\
a^{(4)} &= a_{200} + a_{110} + a_{101} \\
a^{(6)} &= a_{300} + a_{210} + a_{201} + a_{120} + a_{102} + a_{111} \\
a^{(8)} &= a_{400} + a_{310} + a_{301} + a_{220} + a_{202} + a_{211} + a_{130} + a_{103} + a_{121} + a_{112} \\
a^{(10)} &= a_{500} + a_{410} + a_{401} + a_{320} + a_{302} + a_{311} + a_{230} + a_{203} + a_{221} + a_{212} \\
&\quad + a_{140} + a_{104} + a_{131} + a_{113} + a_{122}
\end{aligned} \tag{1.26}$$

Both $L^{(2m)}$ and $M^{(2m)}$ can be evaluated in terms of N -photon propagators.

1. To proceed, one needs something like a Bern-Kosower [19] type master formula for the electron line dressed with any number of photons, with a single constant external (arbitrarily weak) magnetic field insertion. For the magnetic moment calculation, the external vertex is distinguished by its $\sigma^{\mu\nu} = \frac{1}{2}[\gamma^\mu, \gamma^\nu]$ form (1.3), while all internal, virtual photon vertices are of the usual γ^μ form.
2. Please write down the worldline formula for the anomalous magnetic moment of the electron $a = \tilde{F}_2(0)$, corresponding to Dirac trace expression (1.5) for M .
3. As the external vertex transfers a (vanishing) momentum, the incoming and outgoing electron on-mass shell legs are distinct, and thus there are three kinds of N -photon propagators; k photons crossing the external vertex (cross-photons) and $m[m']$ photons originating and terminating on the incoming [outgoing] electron leg (leg-photons). One needs to prove in the worldline formalism that each $km'm$ integral (corresponding to a set of quenched set of 1-particle irreducible Feynman vertex diagrams without electron loops) is separately (i) a gauge-invariant set, and (ii) the minimal gauge invariant set.
4. Hopefully the distinction motivates the gauge set sign rule (1.11). Keep in mind, however, that this empirical rule is already violated by the gauge set (2, 2, 0).
5. Please write down the worldline integral for one-loop anomaly $a_0^{(2)}$ in (1.7). The first thing to verify is that the worldline (1, 0, 0) integral reproduces Schwinger's $\frac{1}{2} \left(\frac{\alpha}{\pi}\right)$ result [156], exactly. That is an exercise in converting the integral into Feynman-parametric form, already done several times for other amplitudes.
6. Please write down the worldline integral for 2-loop anomaly $a_0^{(4)} = M^{(4)} - L^{(2)}M^{(2)}$ in (1.7). Can $L^{(2)}M^{(2)}$ be absorbed into the integrand? If cancelations can be made pointwise, that would obviate a need for constructing UV (and IR?) counterterms.
7. For 2-loop anomaly there are only 2 quenched gauge sets $km'm$: (2, 0, 0) and (1, 1, 0), which equals (1, 0, 1) by time reversal, see figure 1.2 and figure 1.5. So, reformulate the 2-loop calculation as two worldline integrals, one for each gauge set. Most likely, want to do the gauge set (2, 0, 0) first, as it seems to have simpler subdiagram structure (though not sure about that). Do not attempt (for

now) to evaluate these analytically (though Broadhurst, Laporta, Kreimer, etc., would be interested to see whether some simplification occurs), main thing is to understand that the UV renormalization works, and that there are no intermediate IR divergences in this reformulation.

1.4 Volkov method

In ref. [178] Volkov explains that $A_1^{(2n)}$ is free from infrared divergences since they are removed by the on-shell renormalization. However, Volkov also states that there is no universal method in QED for canceling IR divergences in the Feynman graphs analogous to the R operation, and that the standard subtractive on-shell renormalization cannot remove IR divergences point-by-point in Feynman-parametric space, as it does for UV divergences. Moreover, it can generate additional IR-divergences.

That QED on-mass shell amplitudes are IR-free must be an old result; even I have several papers generalizing that to QCD [39, 40, 42, 51]. Tom Kinoshita and I solved the problem of point-by-point removal of IR divergences in Feynman-parametric space in my thesis [49], with a super-elegant formula (who needs forests?) for the UV and IR finite part of amplitude M_G ,

$$\Delta M_G = \prod_{ij} (1 - I_{G/S_i})(1 - K_{G/S_j}) M_G, \quad (1.27)$$

where the products are over all self-energy and vertex subdiagrams S_i and S_j . I have a bright memory of figuring out how to do it one quiet evening in Ithaca, babysitting for a friend's toddler. But, as Volkov [178] and Aoyama *et al.* [10] explain, our approach was apparently not general enough to deal with the 4- and 5-loop contributions.

Volkov's algorithm is developed in *New method of computing the contributions of graphs without lepton loops to the electron anomalous magnetic moment in QED* [179]. It is based on the ideas used for proving UV-finiteness of renormalized Feynman amplitudes [5, 168]. He focuses on n -loop graphs with no lepton loops, or, in the notation of these notes, $a^{(2n)}[V]$. Volkov calculation groups Feynman graphs by self-energy graphs families because they have similar integrand structure. In contrast to refs. [9, 31, 32, 50] he does not evaluate these self-energy graphs directly; all his calculations are performed with vertex graphs, i.e., precisely what is needed to evaluate gauge sets of sect. 1.2. However, as illustrated in figure 1.4, each gauge-set vertex diagram belongs to a different self-energy diagram, so Volkov calculation will require a major reorganization of how integrands are generated, requiring months of recoding.

So far Volkov has evaluated the ladder graph and the fully crossed graph up to 5 loops. The cross graphs are of interest because they do not contain divergent subgraphs, so their contributions only depend on the gauge, but not on the choice of subtraction procedure.

While the contributions of individual vertex graphs (and self-energy sets [9]) are all over the place, all gauge-invariant sets are insanely small up to order 8, and it would be very sweet to see that this continues through order 10 (at least for the 5-loop graphs with no electron loops). My hunch is that starting with the gauge set (5, 0, 0) of figure 1.5 (5! vertex graphs, some of them symmetric pairs) would be the most rewarding. Stefano

Laporta thinks it too hard, and suggests starting with the 5-loop relative $(1, 3, 1)$ (or $(1, 2, 2)$) of the 4-loop set $(1, 2, 1)$, which would entail less than $5!$ vertex graphs (I have not counted how many). As no high accuracy is needed, a numerical check of the QED finiteness conjecture would good enough if the gauge sets evaluated to two significant digits or so, but even that will need a lot of computer time.

1.5 Hopf algebraic approach

Hopf algebraic approach of Kreimer and collaborators [29, 107, 114, 115] is very appealing - it is just that I personally have no clue how to turn it into a direct $(g - 2)$ gauge set calculation. In the 2008 paper [115] Dirk Kreimer and Karen Yeats write:

“One case where there is a natural interpretation is QED with a linear number of generators, namely

$$X_1 = 1 + \sum_{k \geq 1} p(k) x^k \frac{X_1^{2k+1}}{(1 - X_2)^{2k} (1 - X_3)^{2k}}, \quad (1.28)$$

with X_2 and X_3 as before and with $p(k)$ linear, which corresponds to counting with Cvitanović’s gauge invariant sectors [41].”

Even in this simple case I do not see how this counts the gauge sets. My generating function for G_{2n} , the number of gauge sets (eq. (7) in ref. [41]) is

$$\sum_{n=1}^{\infty} G_{2n} = \frac{X}{(1 + X)(1 - X)^3}. \quad (1.29)$$

Broadhurst, Delbourgo and Kreimer [29] 1996 *Unknotting the polarized vacuum of quenched QED* unearthes much knot-theory magic, leading to cancelations of “transcendentals.” While their particular conjecture did not work out in higher orders, the conceptual scheme might be another route to proving the QED is finite - if there is some finite knot-theory basis for expressing the value of every gauge set, and the gauge invariance induced cancelations are so strong to lead to the large cancelations of transcendentals (hyperlogarithms), then perhaps that gives bounds on the size of each gauge set which are slower than combinatorial. The number of different kinds of knots with n crossings is known to grow only exponentially, not faster.

Henry Kibler (on page 52) has a fresh idea for how to approach the finiteness conjecture, using the *Hepp bound*, see Panzer 2018-06-07 below.

Note that the Kibler and Kreimer [107] definition of a “gauge set” differs from (1.10) used here. They organize a gauge-dependent calculation into “gauge sets” of different parameter dependence.

My notes on these papers are below, starting on page 49.

1.6 Method of smooth conjugacies

If Feynman knew Poincaré: How to replace many diagrams by one

In quantum field theory the standard Feynman diagram methods become quickly unwieldy at higher orders. However, it is frequently observed that the sums of Feynman diagrams, each individually complicated, simplify miraculously to rather compact expressions.

Here comes a possible reason why that can be traced back to Poincaré, and is perhaps not something that a field theorist would instinctively hark to as a method of computing perturbative corrections: make the dynamics linear (“free”) by flattening out the vicinity of a path integral extremum by a smooth nonlinear coordinate transformation. The resulting perturbative expansion is more compact than the standard Feynman diagram perturbation theory.

The smooth conjugacy method sketched here would require some serious work to make it a workable quantum field theory scheme. The reader might prefer to skip straight to the worldline formalism sect. 1.3.

The periodic orbit theory is a classical, deterministic theory [45] that describes nonlinear systems in “chaotic” (for low-dimensional systems) or “turbulent” (for PDEs) regimes. The theory allows us to calculate long time averages in a chaotic system as expansions in terms of the periodic orbits (cycles) of the system. The simplest example (the deterministic analogue of the quantum evolution operator) is provided by the Perron-Frobenius operator

$$\mathcal{L}\rho(x') = \int dx \delta(f(x) - x')\rho(x) \quad (1.30)$$

for a *deterministic* map $f(x)$ which maps a density distribution $\rho(x)$ forward one integer step in time. The periodic orbit theory relates the spectrum of this operator and its weighted evolution operator generalizations to the periodic orbits via trace formulas, dynamical zeta functions and spectral determinants [45, 79].

For quantum mechanics the periodic orbit theory is exact on the semiclassical level [85], whereas the quintessentially quantum effects such as creeping, tunneling and diffraction have to be included as corrections. In particular, the higher order \hbar corrections can be computed perturbatively by means of Feynman diagrammatic expansions [79]. We illustrate how this works by the parallel, but simpler example of *stochastic* dynamics. Cvitanović [44] *Chaotic Field Theory: A sketch* is a programmatic statement how this theory might connect to quantum field theory, and, by a way of motivation, an easy introduction into different approaches to incorporating stochastic corrections into classical dynamics.

What motivated the work [46, 47, 52] summarized in ref. [44] is the fact that the form of perturbative corrections for the stochastic problem is the same as for the quantum problem, and still the actual calculations are sufficiently simple that one can explore more orders in perturbation theory than would be possible for a full-fledged quantum theory. For the simple system studied, the result is a stochastic analog of the Gutzwiller trace formula with the “ \hbar corrections” computed to five orders beyond what has been attainable in the quantum-mechanical applications. Already a discrete time, 1-dimensional discrete Langevin equation [98, 119],

$$x_{n+1} = f(x_n) + \sigma\xi_n, \quad (1.31)$$

with ξ_n independent normalized random variables, suffices to reveal the structure of perturbative corrections. We treat a chaotic system with weak external noise by replacing the deterministic evolution δ -function kernel of Perron-Frobenius operator (1.30) by \mathcal{L}_{FP} , the Fokker-Planck kernel corresponding to (1.31), a peaked noise distribution function

$$\mathcal{L}_{FP}(x', x) = \delta_\sigma(f(x) - x'). \quad (1.32)$$

In the weak noise limit the kernel is sharply peaked, so it makes sense to expand it in terms of the Dirac delta function and its derivatives:

$$\delta_\sigma(y) = \sum_{m=0}^{\infty} \frac{a_m \sigma^m}{m!} \delta^{(m)}(y) = \delta(y) + a_2 \frac{\sigma^2}{2} \delta^{(2)}(y) + a_3 \frac{\sigma^3}{6} \delta^{(3)}(y) + \dots \quad (1.33)$$

where

$$\delta^{(k)}(y) = \frac{\partial^k}{\partial y^k} \delta(y),$$

and the coefficients a_m depend on the choice of the kernel. We have omitted the $\delta^{(1)}(y)$ term in the above because in our applications we shall impose the saddle-point condition, that is, we shift x by a constant to ensure that the noise peak corresponds to $y = 0$, so $\delta'_\sigma(0) = 0$. For example, if $\delta_\sigma(y)$ is a Gaussian kernel, it can be expanded as

$$\begin{aligned} \delta_\sigma(y) &= \frac{1}{\sqrt{2\pi\sigma^2}} e^{-y^2/2\sigma^2} = \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{n!2^n} \delta^{(2n)}(y) \\ &= \delta(y) + \frac{\sigma^2}{2} \delta^{(2)}(y) + \frac{\sigma^4}{8} \delta^{(4)}(y) + \dots \end{aligned} \quad (1.34)$$


Analogies between noise and quantum mechanics can be explored by casting stochastic dynamics into path integral form (a stochastic Wiener integral). The periodic orbit theory is a nonperturbative, ‘‘WKB’’ method for approximating such integrals, which can then be improved by systematic perturbative corrections. In the weak noise case the standard perturbation theory is an expansion in terms of Feynman diagrams. For semiclassical quantum mechanics of a classically chaotic system such calculation was first carried out by Gaspard [79]. The stochastic version, implemented by Dettmann *et al.* [46], reveals features not so readily apparent in the quantum calculation. Perhaps some of these could be of interest to Kreimer and collaborators, sect. 1.5.

The Feynman diagram method becomes quickly unwieldy at higher orders.³ However, in the Feynman diagram approach pursued in ref. [46], the authors observe that the sums of Feynman diagrams simplify miraculously to rather compact expressions.

Now the surprise; one can compute the same corrections faster and to a higher order in perturbation theory by integrating over the neighborhood of a given saddlepoint *exactly* by means of a nonlinear change of field variables. This elegant idea of flattening the neighborhood of a saddlepoint, introduced by Mainieri *et al.* [47], and referred to here as the *smooth conjugation method*, is perhaps an altogether new idea in field

³ The matrix method, introduced by Vattay *et al.* [52], based on Rugh’s [144] explicit matrix representation of the evolution operator will not be discussed here. If one is interested in evaluating numerically many orders of perturbation theory and many eigenvalues, this method is unsurpassed.

theory. The idea, that can be traced back to Poincaré [134], injects into field theory a method standard in the construction of normal forms for bifurcations [99]: perform a smooth nonlinear coordinate transformation $x = h(y)$, $f(x) = h(g(h^{-1}(x)))$ that flattens out the vicinity of a fixed point and makes the map *linear* in an open neighborhood, $f(x) \rightarrow g(y) = \mathbf{J} \cdot y$.



The resulting perturbative expansion turns out to be more compact than the standard Feynman diagram perturbation theory; whether it is better than the traditional loop expansions for computing field-theoretic saddlepoint correction remains to be seen.

What is new is that the problem is being solved locally, periodic orbit by periodic orbit, by translation to coordinates intrinsic to the periodic orbit.

This local rectification of a map can be implemented only for isolated non-degenerate fixed points (otherwise higher terms are required by the normal form expansion around the point), and only in finite neighborhoods, as the conjugating functions in general have finite radii of convergence.

In this approach the neighborhood of each saddlepoint is rectified by an appropriate nonlinear field transformation, with the focus shifted from the dynamics in the original field variables to the properties of the conjugacy transformation. The expressions thus obtained *correspond to sums* of Feynman diagrams, but are more compact.

We will try to explain this simplification in geometric terms that might be applicable to more general field theoretic problems. The idea is this: as the dynamics is nonlinear, why not search for a nonlinear field transformation $\phi = h(\tilde{\phi})$ (a smooth conjugacy) that makes the intrinsic coordinates as simple as possible? Schematically –wrong in detail, but right in spirit– find a smooth conjugacy such that the action $S[\phi] = S_0[\phi] + S_I[\phi]$ in the partition function path integral becomes the free, quadratic action,

$$Z = e^W = \int [d\phi] e^{S[\phi]} = \int [d\tilde{\phi}] \frac{1}{|\det \partial h(\tilde{\phi})|^{\frac{1}{2}}} e^{\frac{1}{2} \tilde{\phi}^\top \frac{1}{\Delta} \tilde{\phi}}, \quad (1.35)$$

at the price of having the determinant of the conjugacy Jacobian show up as a weight.

Ref. [46] treats the problem of computing the spectrum of this operator by standard field-theoretic Feynman diagram expansions. Here we formulate the perturbative expansion in terms of smooth conjugacies and recursively evaluated derivatives. The procedure, which is relatively easy to automatize, enables us to go one order further in the perturbation theory, with much less computational effort than Feynman diagrammatic expansions would require.

[TO BE CONTINUED]

1.7 Summary

Everyone makes mistakes—including Feynman

— Toichiro Kinoshita [102]

Currently Sergey A. Volkov is in the best position to check the QED finiteness conjecture numerically, by computing the 5-loops gauge sets.

Worldline formalism could be useful on a qualitative level, as a way of proving the finiteness of QED conjecture,

1. Develop a saddle point expansion for the N -photon propagator integrals, such that the leading term explains the apparent $\approx \pm 1/2$ (or a multiple thereof) size of each quenched gauge set. Affleck *et al.* [1] and G. Torgrimsson *et al.* [175] show the way.
2. Use that to establish bounds on gauge sets for large orders, prove finiteness of quenched QED. If that works, I trust electron loop insertions will be next, and thereafter renormalons [121], etc., will go **gently into that good night**.

and in a precise way, as a new computational tool:

1. Develop a worldline formulation of spinor QED in which each gauge set is given by a computable integral, in a way to be fleshed out in sect. 1.3.2.
2. Parenthetically, a reformulation of the self-energy diagrams magnetic moment calculation, sect. 1.2.3, would be an even greater computational time saver - all quenched diagrams contributions calculated at one go.
3. In either case, a worldline formulation might make it possible to evaluate orders beyond 5-loops, as the number of gauge sets grows only polynomially. A win-win.
4. A gauge set is by definition UV and IR finite. The worldline formalism quenched QED needs wave function counterterms, as in (1.6).
5. Things get interesting with reformulating the quenched 3-loop calculation as four worldline integrals / gauge sets, see figure 1.4 and figure 1.5. In particular, the fermion line attachments of different kinds of N -photon propagators now get intertwined.
6. One electron-loop insertion into $(1, 0, 0)$ might be the easiest worldline integral to evaluate, but I find the quenched sets a higher priority.
7. One photon-photon scattering electron-loop insertion into $(2, 0, 0)$ might be the most tempting to evaluate, but I find the quenched sets a higher priority.

My main problem at the moment (well, there are many:) is that nobody seems to have written an explicit formula for the spinor QED anomalous magnetic moment in the worldline formalism.

1.8 Is QED finite? Correspondence

I am not sure I'll will tell you anything useful any time soon, as I'm a lapsed field theorist, but here is for your amusement my [recent rant](#). To cite my wife: "You may well be right".

— [Keep the Aspidistra Flying!](#)

2007-03-18 Dirk Kreimer I was pretty much looking forward to grab you for a chat on Dyson Schwinger etc whilst in Bonn this starting week. Now I fail to be there myself due to a stupid mechanical problem with a vertebrae. Pondering about other options, let me just point out that the IHES (www.ihes.fr) has an excellent visitor program. If you foresee some time you fancy to pay us a visit, please go ahead.

I am running a little seminar at Boston whilst being there, so besides what works out with IHES, whenever you are available in the above period, you are herewith invited for a talk at BU. Would you be able to come for a seminar to Boston, Thu, Nov 29 this year?

2011-08-23 Predrag to Dirk:

I feel like some stuck-up god knows who for not having answered sooner - but it is not that. I have been focusing on turbulence (with the goal of using what I learn on Yang-Mills, eventually - if it works, Feynman diagrammar will be of secondary importance, the first step is fully and totally non-perturbative) so I feel like fraud talking about QFT when I am suspended between the perturbative past and the elusive non-perturbative future.

If someone wants to hear about turbulence as in ChaosBook.org/tutorials, I can do that honestly - otherwise we wait until I actually do any QFT worth hearing about...

I got birdtracks.eu published, finally - you get 100 birdtracks per 1\$, a real deal if you are into that kind of thing.

Guilty as Charged - Predrag

2011-08-26 Dirk June 01-05 2009 I am organizing a workshop at IHES. It has a non-perturbative component via Dyson-Schwinger eqs. Would you be interested to participate and come?

2017-06-16 Dirk I will show your notes to Henry Kissler [[106](#)] <kissler@physik.hu-berlin.de> and [Michael Borinsky](#) [[25](#), [26](#)] <borinsky@physik.hu-berlin.de> .

There should be a few people at Les Houches June 2018 who have something to say about asymptotics of solutions of Dyson-Schwinger equation for example.

2013-10-23 Warren D. Smith warren.wds@gmail.com

(Predrag: I do not know Dr. Smith, but for his life until 2009, see his resume [here](#).)

I found out interesting info which the textbooks do not know!

Cvitanović 1977 conjectured that gauge-invariant quenched diagram sets always have small sum of diagram values, indeed small enough that he thought quenched QED series would always converge [41]⁴. He apparently originally thought this even for unquenched but Lautrup [121] disproved it and Dyson had long had a highly convincing argument [61] (Reprinted p.255-6 in Selected papers of Freeman Dyson with commentary, AMS 1996) – and I now have even more convincing arguments – that generic QED series diverge for any nonzero α .) This was due to Cvitanović’s empirical observation of amazingly huge cancellations within gauge-invariant diagram sets, especially in quenched QED.

However, what Cvitanović did not know, was that **Bogomolny** and Kubyshin 1981-1982 found estimates of the growth rate of generic QED series, and also for the quenched QED subseries, and indeed for QED for diagrams with k electron loops only (k fixed) [22]. I only found this out the other day, but I had long known about estimates due to Dyson and others predicting divergence for QED series. It is just that this B+K work by permitting arbitrary fixed k , directly addresses Cvitanović’s quenched-convergence conjecture and massively conflicts with it – for quenched QED the prediction is that at N th order we get a quenched diagram sum growing factorially with N :

Evgeny B Bogomolny and Yu A Kubyshin:

1. The choice of the form of the steepest-descent solutions [22]. Asymptotic estimates for diagrams with a fixed number of fermion loops in quantum electrodynamics.
2. The extremal configurations with the symmetry group $O(2) \times O(3)$, Soviet J. Nuclear Phys. 35 ,1 (1982) 114-119.⁵

I believe in the sort of arguments Bo+Ku are making, albeit the details are questionable. (E.g. saddlepoint asymptotic estimates are not rigorous unless you prove stuff about the saddlepoints and about tail estimates, which they never proved, and probably nobody can prove.)

Note that this believed $N!$ growth for fixed- k (including quenched) QED diagrams (Lautrup’s diagrams also feature $N!$ growth) is far faster than the believed growth – more like $(N/2)!$ – for the *full* QED series!! This fact that subseries diverge far more rapidly than full series can only be explained by presuming that the values at different k cancel each other amazingly well when we sum over all k . This indicates Cvitanović was extremely wrong asymptotically, and that cancellations quite different than his observations ultimately occur which seem even more dramatic.

However, Cvitanović was correct that amazingly large cancellations (also) occur, empirically, within the quenched diagrams alone, at least for the small N that have been reached by computer. And I presume that Kreiman’s knot ideas [114] and my “fractal distribution” empirical observations are a partial explanation of why.

⁴Predrag: ‘quenched’ = no internal electron loops, i.e., $m_e \rightarrow \infty$ approximation

⁵Predrag: there is a hard copy at GT library, 4th Floor East Call Number: QC173.J252X (or microfilm?)

So a consistent picture is now developing about how QED perturbative series (and various interesting sub-series, such as quenched) allegedly behave asymptotically (Although I never saw anything saying all that I just said in one place...)

2013-10-23 Warren to Predrag do you have a graph-theoretic characterization of ‘gauge invariant diagram set’ or know how many such (minimal) sets there are at order N ? E.g. does their count grow exponentially, super-exponentially, polynomially or what? You gave a formula for the count of quenched gauge sets which grows only polynomially but I suspect exponential or faster growth for unquenched QED. Even an incomplete graph-theory understanding might be adequate to get good growth bounds.

2013-10-23 Warren Although I do not understand “gauge invariant sets of Feynman diagrams” I have figured out enough now to prove that their count grows ultimately superexponentially in unquenched QED. Specifically I now claim to have a proof that the number of gauge invariant sets of Feynman diagrams in QED at α^n order is

$$\approx 0.01 (96)^{-n/4} \frac{n!}{(n/2)!(n/4)!}$$

for any integers $n > 0$. And this clearly grows superexponentially. (My bound is very unlikely to be optimal.)

2013-10-23 Predrag to Warren : I’m interested in this discussion, but can you do me a favor and actually read my paper, and edit your initial email accordingly, before we wrangle with further details?

Many of your statements are addressed in my paper, and I can answer them more efficiently if you go through them first. I love Dyson dearly, but his statement is an elementary statement about asymptotic series for factorials. Mine is about mass-shell gauge invariant quantities (perhaps planar?), a topic that is gaining some traction now, unfortunately not applicable to QED.

If you can get gauge sets added up, that would be useful, but Aoyama *et al.* [7, 9, 11] do not have them: they use the second method of computing magnetic moment, eq (6.22) (see sect. 1.2.3 above). That mixes up the gauge sets.

2013-11-23 Warren Smith [draft of my notes](#) .

2013-10-24 Warren Predrag: Bogomolny and Kubyshin [22] should find combinatorial growth in all sectors - a perturbation expansion around a saddle point is always asymptotic. For example, a path integral over a non-linear oscillator with quartic potential can be well defined as integral, but saddle point expansion is asymptotic - a simple example is worked out in Sect 3.3 *Saddle-point expansions are asymptotic* ([click here](#)).

I agree with Dyson (and Bogomolny & Y. Kubyshin, which is more modern version of the argument).

Essentially, what they did is this. “Instanton” solutions of quantum field theory stationarize the action and hence are “saddlepoints.” The full path integral over

all quantum field histories, is hopefully dominated by this saddlepoint and nearby configurations. It seems there are more than one kind of instanton, and you need to know which one is the “dominant” saddlepoint, and they may have the wrong one, but if so it hopefully does not matter much (changes some constants, but not the qualitative behavior, of their results, one hopes). So anyhow, under this assumption they actually are able to work out the asymptotics of the Nth term in QFT perturbative series, in the limit where N is large. They find $N!^P$ style growth for positive powers P. Furthermore, for the “quenched” QED sub-series, one might naively think that approach would say nothing about it, but they have further generating functionology tricks which enables them to get conjectured asymptotics for that, and lots of other sub-series, too.

The result is $N!^P$ style growth in all cases. Indicating divergency and with radius of convergence zero.

Suslov [170] and other Russians claim the original idea for this was due to Lev Lipatov, but it has been explored by a fairly large number of papers & authors now, I think mostly French & Russian. My tome has 2 chapters 5 & 6 on Dysonian & other divergence arguments which give pointers into the literature. The Suslov paper you cite also has many such pointers.

Predrag: What I claim/hope is that gauge invariance + mass-shell condition (neither accounted for in the above asymptotic estimates) induce cancellations that make the theory convergent.

–I think this literature already had invented some way of handling gauge invariance within their saddle pointage. They had thought of it and figured out a way to deal with it. (Mind you, all of this stuff is horribly nonrigorous.) And if by “mass shell” you mean what I am calling “quenched QED” (kind of a battle over which name is worse...) then as I said, Bogol already had a trick for obtaining that from the unquenched analysis.

Predrag: Mass shell means that all external legs of Feynman diagrams are the physical, asymptotic states satisfying $E = mc^2$. Intermediate virtual states do not do that. Gauge invariance cancellations kick in for the mass shell states, not for the off-mass shell amplitudes. Saddle point estimates do not use mass-shell conditions, to the best of my knowledge.

Suslov [170] *High orders of perturbation theory. Are renormalons significant?*, [arXiv:hep-ph/0002051](https://arxiv.org/abs/hep-ph/0002051), seems to be about an unphysical but comparatively well behaved ϕ^4 quantum field theory. It seems to argue that renormalons of t’Hooft and Lautrup do not matter. I have not studied Suslov’s papers (or the ones that cite them inspirehep.net/record/510344/citations).

The Lautrup renormalon does not contradict the instanton-based results. And renormalons also lead to yet more kinds of series divergency. By the way my tome also discusses Lautrup renormalon. Suslov in his footnote 4 and nearby

points out that the whole saddlepoint approach of Lipatov is nonrigorous (like I didn't know) and in fact may be bogus. Physicists often use saddlepoint analysis but rarely do what it takes to do it rigorously, for example they rarely prove "tail estimates." (And of course when everything is in an infinite dimensional space like it is here, that adds a whole new level of difficulty in trying to get any rigor.) They generally instead just hope for the best. Suslov suggests that the renormalon may be a clue that in this case, the tail may be big enough that the whole Lipatov technique is wrong.

I also think everything Suslov does re Borel transforms, is wrong for QED since QED is not going to be Borel summable. However for ϕ^4 theory Borel might be OK... yes, it is, see table 11 in my tome.

Predrag: You have to learn all this stuff (including [planar field theory](#)) to get people to read you.

I don't know whether anybody will read me, and I've had plenty of trouble learning what I did learn, and I'm not willing to do too much work to overcome that. I am certainly quite ignorant in many respects. I draw some comfort from the fact that many leaders in this area, including many Nobel prize winners, have made plenty of errors, some of which I detected for the first time, and some of which were real howlers. The moral of that for me is, nobody really knows what they are doing about this stuff. You've got to know the right stuff, not all stuff. I hope if my ideas are valid they will eventually gain some attention. If they are wrong I hope this will become clear as soon as possible...

2017-04-30 Predrag to Stefano Laporta <stefano.laporta@bo.infn.it>:

Thanks for the listings of gauge sets, and of course, for your whole amazing project. While in detail my 1977 guesses are wrong, the overall finiteness conjecture still looks good - it's crazy how small all these individual contributions are.

The reason I got into a collaboration with Kinoshita (nominally, Tung-Mow Yan was my adviser) is that he gave me the 3-loop ladder diagram (at 4-loops, it would belong to your gauge set $(6) = (4, 0, 0)$) to evaluate, and I derived a compact formula for the parametric integrand of n -loops ladder diagram. So maybe gauge sets of type $(6) = (4, 0, 0)$ are the easiest to evaluate?

Is evaluating gauge set $(5, 0, 0)$ with 5 loops feasible? If that turns out to be $+1/2$, that would be sweet.

2017-05-05 Laporta to Predrag

To my knowledge, there are no ideas how to bound the sizes of the gauge sets.

In the set $(6) = (4, 0, 0)$ the ladder is easy to evaluate, but the others are not. The evaluation of the set $(5, 0, 0)$ with five loops would be a real challenge.

Among the sets (1)-(6), the easiest to evaluate is the set $(3) = (1, 2, 1)$.




Figure 1.8: Examples of the $A_1^{(8)}$ 4-loop vertex diagrams belonging to the 25 gauge-invariant sets. The number indicates the gauge-invariant set to which the diagram belongs. Gauge sets (1) = (1, 3, 0), (2) = (2, 2, 0), (3) = (1, 2, 1), (4) = (2, 1, 1), (5) = (3, 1, 0), (6) = (4, 0, 0). In the case of the sets 1-16, 24 and 25, remaining diagrams in the set can be obtained by permuting separately the vertices on the left and right side of the main electron line, and considering also the mirror images of the diagrams. From Laporta [116].

| | | | |
|----|---------|---|---------|
| 1 | (1,3,0) | - 1.971075616835818943645699655337264406980 | - 1/2 |
| 2 | (2,2,0) | - 0.142487379799872157235945291684857370994 | 1/2 (!) |
| 3 | (1,2,1) | - 0.621921063535072522104091223479317643540 | - 1/2 |
| 4 | (2,1,1) | 1.086698394475818687601961404690600972373 | 1/2 |
| 5 | (3,1,0) | - 1.040542410012582012539438620994249955094 | - 1/2 |
| 6 | (4,0,0) | 0.512462047967986870479954030909194465565 | 1/2 |
| 7 | | 0.690448347591261501528101600354802517732 | |
| 8 | | - 0.056336090170533315910959439910250595939 | |
| 9 | | 0.409217028479188586590553833614638435425 | |
| 10 | | 0.374357934811899949081953855414943578759 | |
| 11 | | - 0.091305840068696773426479566945788826481 | |
| 12 | | 0.017853686549808578110691748056565649168 | |
| 13 | | - 0.034179376078562729210191880996726218580 | |
| 14 | | 0.006504148381814640990365761897425802288 | |
| 15 | | - 0.572471862194781916152750849945181037311 | |
| 16 | | 0.151989599685819639625280516106513042070 | |
| 17 | | 0.000876865858889990697913748939713726165 | |
| 18 | | 0.015325282902013380844497471345160318673 | |
| 19 | | 0.011130913987517388830956500920570148123 | |
| 20 | | 0.049513202559526235110472234651204851710 | |
| 21 | | - 1.138822876459974505563154431181111707424 | |
| 22 | | 0.598842072031421820464649513201747727836 | |
| 23 | | 0.822284485811034346719894048799598422606 | |
| 24 | | - 0.872657392077131517978401982381415610384 | |
| 25 | | - 0.117949868787420797062780493486346339829 | |

Table 1.2: Contribution to $A_1^{(8)}$ of the 25 gauge-invariant sets of figure 1.8, as reported by Laporta [116].

2017-06-17 Laporta on *transcendentals*: There are no cancellations of transcendentals at 2 loops.

At 3 loops there is only one:

$\zeta(2) \log(2)$ cancels in the sum of the gauge sets (1,2,0)+(2,1,0)+(1,1,1)

At 4 loop there are several:

$\zeta(2)^2 \log(2)$ cancels in the sum of the gauge sets 8,9,10 (using my numbering of gauge sets)

$\zeta(2)^2$ cancels in the sum of the gauge sets 5,6,20 (or 10,13,19)

a weight-7 elliptic constant cancels in the sum of the gauge sets 2,3,4; (I have not checked all possible combinations of gauge sets)

2017-06-16 Predrag to Stefano thanks for the info! Apparently there are not so many transcendentals cancellations when one is adding gauge sets...

I was actually thinking of cancellations *within* an individual gauge set. For example, in Laporta and Remiddi [118] *The analytical value of the electron (g-2) at order in QED* you list analytical values for each diagram separately + $\ln \lambda$

terms. They have $\zeta(2) \log(2)$'s and $\zeta(2) \log(2)^2$ and zillion other transcendentals - do many of them cancel within a gauge set (for example, $\ln \lambda$'s certainly do)?

Just trying to figure out if we can use Kreimer kind of ideas to get any bounds.

2017-05-25 Predrag to Stefano:

In the literature there seem to be 2 approaches that might be relevant to establishing bounds on, and perhaps even to actual computation of gauge-invariant sets (ignoring various $N = 2$ and $N = 4$ supersymmetric models), the worldline formalism pursued by Schubert and collaborators, see sect. 1.3, and Hopf algebraic approach of Kreimer and collaborators, see sect. 1.5. The worldline formalism operates with N -photon propagators, the ingredient that defines the quenched gauge-invariant sets for $(g - 2)$. The summary of this literature is in sect. 1.2.4.

2017-05-21 Predrag to Sergey A. Volkov <volkoff_sergey@mail.ru>

I have read your *New method of computing the contributions of graphs without lepton loops to the electron anomalous magnetic moment in QED* [179] (and the earlier ref. [178]) with great interest, and I can see that you are set to compute the 5-loop correction to the electron $(g - 2)$. This is a very hard calculation, and approaching it strategically is a necessity.

May I suggest that you order your calculation by gauge sets of figure 1.5, as illustrated (for the 4-loop case) in Stefano Laporta's figure 1.6? My hunch is that the gauge set $(6) = (4, 0, 0)$ would be the most interesting, though Stefano thinks it too hard, and suggests starting with a 5-loop relative $(1, 3, 1)$ (or $(1, 2, 2)$?) of the set $(3) = (1, 2, 1)$ instead. While the contributions of individual vertex graphs (and self-energy sets [9]) are all over the place, all gauge-invariant sets are insanely small up to order 8, and it would be very sweet to see that this continues through order 10 (at least for the 5-loop graphs with no electron loops).

By the way, to check my conjecture one needs the gauge sets only to two significant digits or so, no high accuracy is needed.

2017-06-18 Sergey About gauge sets: It is possible to calculate the contributions of $km'm$ by my method. It will require more computer resources than the calculation of the total contribution without splitting into sets, because the sets $km'm$ cut the families that I use for calculation into many pieces. I will try to calculate 5-loop contributions of $km'm$ after some refinement of my code (this requires some months). Even a not precise calculation needs a lot of computer time.

2017-06-18 Sergey About "That QED on-mass shell amplitudes are IR-free must be an old result...": Is there a rigorous mathematical proof that they are IR-free? I didn't find such a proof. I saw only some approximate considerations about this. The proof of the Bogoliubov-Parasiuk theorem is quite large and complicated. I think that the total proof of IR and UV cancellation should be larger. However, all considerations that I saw are much simpler. I think that an existence of such a proof should affect computational methods. However, at the present moment, the best calculation of the QED contributions to $g-2$ (Aoyama, Kinoshita et al.) uses the subtraction procedure that needs a handwork at 5-loop

level (see the 2008 Aoyama *et al.* article [10], and the link to it in the end of their 2012 article [8]). On the other hand, all mathematical proofs of IR cancellations that I saw are connected with IR divergences of other nature (not on-shell). Is there a mathematical proof that the QED contributions to $g - 2$ are finite at any order of the perturbation series (only for each term, not for the sum)?

2017-06-18 Predrag to Sergey I thought that my thesis argument on IR divergences subtraction in the Feynman-parametric space in [49] establishes IR finiteness for $(g-2)$. We show that the UV and IR finite part of amplitude M_G is given by

$$\Delta M_G = \prod_{ij} (1 - I_{G/S_i})(1 - K_{G/S_j}) M_G \quad (1.36)$$

where the products are over all self-energy and vertex subdiagrams S_i and S_j . We then use Ward identities [180] to show that UV and IR divergent parts of counterterms $I_{G/S_i} M_G$, $K_{G/S_j} M_G$, etc. cancel on mass shell (and the finite parts contribute to the anomaly). In the paper I check various cases, and I seem to have missed some that do not arise in the 3-loop order. But how could anything be simpler than (1.36)? And how could such elegant formula be wrong?

I think the UV/IR finiteness is very clean and clear to see in the dimensional regularization. I have now reread my 1976 *Yang-Mills theory of the mass-shell* [40], and it is a rather amusing, but probably not a persuasive answer to your query. The paper is good and original, so it is never cited. At least it got published. The original draft was to be exactly *one* page of Physical Review Letters, had *one* reference, a homage to Schwinger [156]. Those were happier times - I wrote it in a couple of days while visiting Helsinki Research Institute for Theoretical Physics, mailed it, impressing my hosts infinitely - they have never seen a PR Lett written at that speed, and *bicycled off into Lapland*. But referees made me quadruple the length of the Letter, making it as boring as any other PR Lett. For the proof of absence of on-mass shell $(g - 2)$ IR divergences I cite no less than three of my own unpublished papers (!!!), and nobody else. What happened? I had understood that (the finiteness of QED conjecture had much to do with it) that doing more Feynman graphs would be stupid (expanding around the wrong vacuum), and thus started my nonlinear detour. I started thinking about functional equations on May 1, 1976, and wrote down *the period doubling fixed point function equation* on May 3, 1976. In my office I do have a very thick folder marked UNPUBLISHED, and it must contain the drafts of these IR papers. But how would I know that anyone would care 40 years later?

The one that has IAS number was actually published [39], see [here](#), and it says:

“The anomaly, expressed in terms of the unrenormalized coupling constant, is free of infrared divergences for QED (i.e., all $a^{(2^n)}$ are IR finite). This is particularly easy to demonstrate using the method for separation of IR divergences given in ref. [49]. The analysis is rather technical, but the result is very intuitive: a Feynman integral M_G contributing to the magnetic moment is IR divergent whenever the

corresponding diagram G can be split into a vertex subdiagram S and a cloud of soft gluons attached to the external quark lines (diagram G/S). In general some gluons in G/S can be hard, provided they are within UV divergent subdiagrams. In all cases, the IR divergent part of M_G factors into M_S times the IR divergent part of $L_{G/S}$, where $L_{G/S}$ is the charge form factor computed from the diagram G/S .”

and so on - the paper is focused on QCD, but I believe it establishes IR finiteness of QED on-mass shell amplitudes, though it most likely not as rigorously as you would like it.

Yennie and S. C. Frautschi and H. Suura [182] *The infrared divergence phenomena and high-energy processes* (cited over 800 times!) and Kinoshita [101] *Mass singularities of Feynman amplitudes* say nothing about cancellation of IR divergences by renormalization. Herzog and Ruijl [87] *The R*-operation for Feynman graphs with generic numerators* and Chetyrkin, Tkachov and Smirnov [34, 35, 165] are not of interest, as R*-operation is not about on-shell IR-divergences. (Herzog and Ruijl [87] cite Chetyrkin, Tkachov and Smirnov [34, 35, 165] but no Volkov [178, 179] or Cvitanović [39, 40, 42, 49, 51] papers.)

In ref. [49] we also cited Johnson and Zumino [96] *Gauge dependence of the wave-function renormalization constant in Quantum Electrodynamics*.

An unrelated IR problem: Kinoshita [100] *Note on the infrared catastrophe*, about which he writes [103]: “I developed a method to handle the infrared problem within the perturbative framework of Feynman–Dyson theory [100]. My insight was not deep enough, however, and I overlooked a subtlety in the infrared problem. This became the source of an error in the calculation of radiative correction to the $\mu - e$ decay.”

2017-06-11 Predrag to

Christian Schubert <schubert@ifm.umich.mx>

Stefano Laporta has recently published analytic values of all 4-loop electron magnetic moment gauge sets, and they look very intriguing. Sergey A. Volkov has started evaluating individual 5-loop vertex diagrams and is in position to estimate numerically the size of 5-loop gauge sets. So it might be a good time to make a new attempt to prove/disprove the QED finiteness conjecture. This set of notes is my current best attempt to motivate this effort.

I do not take my 1977 “gauge-set approximation” very literally - its content is only that if individual gauge sets can be bounded to anything growing slower than combinatorially, quenched QED (and hopefully full QED) is a finite theory, not an asymptotic series. But the form of vertex gauge sets is very suggestive; it is defined in terms of N -photon exchanges. So to me the wordline formalism seem the most promising way forward.

While the quenched QED magnetic moment of electron is the cleanest possible physical calculation one can do, for sociological reasons most effort on high order estimates has gone in other directions. Do you have a formula for spinor QED magnetic moment that one could attempt to analyze more closely?

2017-06-12 Christian I have been fascinated by the finiteness conjecture ever after Gerald Dunne and I were led there, too, from Euler-Heisenberg, Borel analysis, Ritus mass shift and worldline instantons fifteen years ago. The things have picked up a lot just during the last year or so, namely:

1. Idrish Huet, Michel Rausch and I are working on the Affleck, Alvarez and Manton (AAM) exponentiation conjecture [1] in 1+1 QED. We had a nice parameter integral representation for the three-loop Euler-Heisenberg for quite a while, but could not get a sufficient number of weak-field expansion coeffs out of it (so far what we got points to AAM not holding at three loops - rather the coeffs go asymptotically BELOW the AAM prediction - but this is very preliminary). Michel was just visiting, and we now got a really nice algorithm (based on the polynomial invariants of the dihedral group) for the analytical calculation of the coeffs from the more difficult (nonplanar) 3-loop EH diagram.
2. If AAM really fails, then presumably the AAM worldline instanton needs refinement at higher loops. My former student Naser Ahmadinia (now postdoc in Korea) has made some progress with this.
3. For many years I am planning to apply the worldline formalism to (g-2), in particular to the important graphs with the light-by-light subdiagram. For this subdiagram I have, since a long time, an off-shell representation that is permutation symmetric, manifestly gauge invariant, without spurious UV poles, and moreover allows one to trivially integrate out the one low-energy photon leg. What held me back was that the various worldline representations that existed hitherto for the electron line were all somewhat cumbersome, and seemed not suitable for high-order calculations. Precisely this problem we have been working on here for the last half year, and things have fallen into place really nicely, we have a Bern-Kosower type master formula for the electron line dressed with any number of photons, in vacuum and in the presence of a constant field, and one of our students is already programming it. I am now definitely trying to assemble a collaboration to attack the QED (g-2), and I anticipate that your notes will be quite useful for motivating my collaborators.

Gies *et al.* [84] have some intriguing numerical results from worldline Monte Carlo (in section 5). Another thing I find quite interesting are Ritus [23] most recent papers [138–140] on the value of the bare fine structure constant - he confirmed to me in an email from last December (at age 90) that he definitely does not think that the bare charge is infinite.

I am appending a summary of our efforts in this line of work [89] which I wrote as a contribution to the 5th Winter Workshop on Non-Perturbative Quantum Field Theory. 22-24 March 2017, Sophia-Antipolis.

2016-12-18 Vladimir I. Ritus <v_i_ritus@mail.ru> to Christian:

Thank you for your note about the papers by Johnson, Baker, Willey (Phys. Rev. 1967, 1969, 1971) and by Adler, Bardeen (Phys. Rev. 1971).⁶ I forgot them while writing my papers [139, 140]. Some months ago I accidentally saw the abstracts of these papers in one of my notebooks with conspectus of the literature. Johnson et al assertion confirms qualitatively my geometric and holographic duality approach to the problem. Both of them deal with photons without self-energy insertions. Have you any objections to this approach?

Nobody believes in infinite bare charge neither with space distribution more singular than $\delta(x)$ nor as Landau pole at final transfer momentum. Please, reed the several first phases in JRLR-paper (changing the "emitted" to "emitting").

I regret that I could not come to Tomsk, where I was at war time 1941-1943 in evacuation.

2017-06-14 Christian to Ritus Just to make sure that I understand your conjecture correctly - are you claiming that $1/4\pi$ is the value of the bare constant in the quenched approximation, or in the full QED?

2017-06-29 Vladimir Ritus The $1/4\pi$ is the value of the bare fine structure constant in the full QED. It follows from duality of 4-dimensional QED and 2-dimensional quantum field theory. The point charge moves along timelike trajectory, hence it has nonzero mass, but the photons emitted to infinity by this charge are massless as the duality connected with them scalar quanta of pairs emitted by the pointlike mirror in (1+1)-space. Constants α_L and α_B are also geometrical. So, this theory is pure geometrical and connected neither with the divergent perturbation theory series, nor with the different methods of their summing. Please, read refs. [139, 140].

2017-06-18 Naser Ahmadiniaz <ahmadiniaz.naser@gmail.com> (currently a postdoctoral fellow at the Center for Relativistic Laser Science (CoReLS), Institute for Basic Science (IBS), Gwangju, South Korea).

My main research interests are in amplitude calculations in QED, QCD and quantum gravity from the worldline formalism. Recently, we have been interested in (off-shell) tree-level amplitudes for QED processes in vacuum as well as in the presence of classical background fields which will also be applied to higher order corrections.

2017-06-20 James P. Edwards <jedwards@ifm.umich.mx>, [homepage](#):

I am currently working with Christian on the worldline approach. In some of our recent work we have been working on more efficient ways to determine g_2 based upon a worldline expression for the dressed spinor propagator (both in vacuum and in a constant background).

2017-06-11 Predrag to Dirk Kreimer <kreimer@physik.hu-berlin.de>

⁶Predrag: 2017-08-04 K. Johnson, M. Baker, and R. Willey, Phys. Rev. 136, B1111 (1964); *ibid.* 163, 1699 (1967); and S. L. Adler and W. A. Bardeen, Phys. Rev. D 4, 3045 (1971) These papers are about the possibility of the existence of an ultraviolet fixed point α_c .

Stefano Laporta has recently published analytic values of all 4-loop electron magnetic moment gauge sets, and they look very intriguing. It might be a good time to make a new attempt to prove/disprove the QED finiteness conjecture. This set of notes is my current best attempt to motivate this effort.

If individual gauge sets can be bounded to anything growing slower than combinatorially, quenched QED (and hopefully full QED) is a finite theory, not an asymptotic series. The form of vertex gauge sets is suggestive of a way forward; it is defined in terms of N -photon exchanges. Hopf algebras might hold the key, but I do not know how to use these ideas.

I would be grateful for any further pointers to the literature. Anybody else I should send these notes to? And I'm finally ready to stand up and be counted - I can fly to Berlin on a short notice if that helps.

2017-06-14 Dirk Kreimer consider coming to [Les Houches — June 4-15, 2018](#) on structures in local quantum field theory.

1.9 QCD gauge sets - a blog

In 1981 Cvitanović *et al.* [51] constructed gauge invariant subsectors in QCD.

2016-12-10 Predrag Penante [133] 2016 On-shell methods for off-shell quantities in $N=4$ Super Yang-Mills: from scattering amplitudes to form factors and the dilatation operator has an up-to-date review of on-shell methods.

2016-12-26 Predrag Read Cruz-Santiago, Kotko and Staśto [38] 2015 *Scattering amplitudes in the light-front formalism*: “The idea is to divide the process into appropriate gauge invariant components. It turns out that the gauge invariant subsets are invariant under cyclic permutations of the external gluons. This decomposition was proposed in works of [58–61] for the tree level amplitudes. A thorough analysis of the relation between color structures and gauge invariance was done in ref. [51]. The color decomposition principle was extended beyond the tree level to loop amplitudes in [63].”

2016-12-26 Predrag Should also read Dixon [55] 1996 *Calculating scattering amplitudes efficiently*.

2017-05-26 Predrag The decomposition of scattering amplitudes into gauge invariant subsets of diagrams is studied by Boos and Ohl [14, 24]. Boos and Ohl [24] *Minimal gauge invariant classes of tree diagrams in gauge theories*, [arXiv:hep-ph/9903357](#) (see [arXiv:hep-ph/9911437](#) and [arXiv:hep-ph/0307057](#) for more detail) is motivated by applications to Standard Model multi-particle diagrams, mostly at the tree level.

Perturbative calculations require an explicit breaking of gauge invariance for technical reasons and the cancellation of unphysical contributions is not manifest in intermediate stages of calculations. The contribution of a particular Feynman diagram to a scattering amplitude depends in the gauge fixing procedure and has

no physical meaning. the identification of partial sums of Feynman diagrams that are gauge invariant by themselves is of great practical importance. Calculation a subset of diagrams that is not gauge invariant has no predictive power, because they depend on unphysical parameters introduced during the gauge fixing.

A *gauge invariance class* is a minimal subset of Feynman diagrams that is independent of the gauge parameter and satisfies the Slavnov-Taylor identities.

The set of diagrams connected by flavor and gauge flips they call *forest*, a set of diagrams connected by gauge flips they call *grove*. They shown that the groves are the minimal gauge invariance classes of tree Feynman diagrams. In unbroken gauge theories, the permutation symmetry of external gauge quantum numbers can be used to subdivide the scattering amplitude corresponding to a grove further into gauge invariant sub-amplitudes.

This (largely uncited) work seems to have no impact on the $(g - 2)$ gauge sets discussed here.

2017-05-27 Predrag Reuschle and Weinzierl [137] *Decomposition of one-loop QCD amplitudes into primitive amplitudes based on shuffle relations* cite our ref. [51]. They say:

QCD calculations organise the computation of the one-loop amplitude as a sum over smaller pieces, called *primitive amplitudes*. The most important features of a primitive amplitude are gauge invariance and a fixed cyclic ordering of the external legs. Primitive amplitudes should not be confused with *partial amplitudes* (also referred to as a *dual amplitude* or a *color-ordered amplitude*), which are the kinematic coefficients of the independent colour structures. The first step in a discussion of perturbative Yang-Mills is the decoupling of color from kinematics,

$$A_{tot} = \sum c_J A_J \quad (1.37)$$

where A_{tot} represents the total amplitude for a scattering process, A_J are all the possible color structures, and A_J are partial amplitudes which depend only on the kinematical data (momenta and polarizations). Partial amplitudes are gauge invariant, but not necessarily cyclic ordered. Partial amplitudes are far simpler to calculate than the full amplitude. There exist linear relations among the partial amplitudes, called Kleiss-Kuijf relations, which reduce the number of linearly independent partial amplitudes to $(n - 2)!$ The leading contributions in an $1/N$ -expansion (with N being the number of colours) are usually cyclic ordered, the sub-leading parts are in general not. The decomposition of the full one-loop amplitude into partial amplitudes is easily derived. However, it is less trivial to find a decomposition of the partial amplitudes into primitive amplitudes.

There are several possible choices for a basis in colour space. A convenient choice is the colour-flow basis [171].

2017-05-27 Predrag Schuster [154] *Color ordering in QCD*: “ We derive color decompositions of arbitrary tree and one-loop QCD amplitudes into color-ordered objects called primitive amplitudes. ”

2017-05-27 Predrag Zeppenfeld [183] *Diagonalization of color factors* (Georgia Tech has no access to this paper)

2017-05-27 Predrag Edison and Naculich [62] *Symmetric-group decomposition of $SU(N)$ group-theory constraints on four-, five-, and six-point color-ordered amplitudes at all loop orders*: “ Color-ordered amplitudes for the scattering of n particles in the adjoint representation of $SU(N)$ gauge theory satisfy constraints that arise from group theory alone. These constraints break into subsets associated with irreducible representations of the symmetric group S_n , which allows them to be presented in a compact and natural way. ”

2017-05-27 Predrag Kol and Shir [109, 110] *Color structures and permutations* has a useful overview of the literature in the introduction, and is a very interesting read overall.

We may permute (or re-label) the external legs in the expression for a color structure and thereby obtain another color structure. This means that the space of color structures is a representation of S_n , the group of permutations. A natural question is to characterize this representation including its character and its decomposition into irreducible representations (irreps).

The decomposition of color structures into irreps was suggested by Zeppenfeld [183].

The space of tree-level color structures TCS_n of dimension

$$\dim(TCS_n) = (n - 2)! \tag{1.38}$$

is the vector space generated by all diagrams with n external legs and an oriented cubic vertex, which are connected and without loops (trees), where diagrams which differ by the Jacobi identity are to be identified.

The f -based and t -based color structures are related by celebrated Kleiss-Kuijf [108] relations (rederived in this paper).

The original problem, that of capturing symmetries of the partial amplitudes which originate with those of the color structures, is now formulated as the problem of obtaining the S_n character of the space of color structures. It turns out that (at least at tree level) this problem was fully solved in the mathematics literature by Getzler and M. M. Kapranov [81].

The free Lie algebra over some set A , denoted by $L(A)$, is the Lie algebra generated by A with no further relations apart for antisymmetry and the Jacobi identity which are mandated by definition.

Self duality under Young conjugation: for some n values TCS_n is self-dual under Young conjugation, namely under the interchange of rows and columns in the Young diagrams

2017-05-27 Predrag Getzler and Kapranov [81] *Modular operads*: “ We develop a ‘higher genus’ analogue of operads, which we call modular operads, in which graphs replace trees in the definition. We study a functor F on the category of

modular operads, the Feynman transform, which generalizes Kontsevich's graph complexes and also the bar construction for operads. We calculate the Euler characteristic of the Feynman transform, using the theory of symmetric functions: our formula is modelled on Wick's theorem. ”

2017-05-27 Predrag Maltoni *et al.* [125] *Color-flow decomposition of QCD amplitudes*

The *color-flow* decomposition is based on treating the $SU(N)$ gluon field as an $N \times N$ matrix. (PC: I think that is what I actually do.) It has several nice features. First, a similar decomposition exists for all multiparton amplitudes, like the fundamental-representation decomposition. Second, the color-flow decomposition allows for a very efficient calculation of multiparton amplitudes. Third, it is a very natural way to decompose a QCD amplitude. As the name suggests, it is based on the flow of color, so the decomposition has a simple physical interpretation.

To calculate the amplitude, one orders the gluons clockwise, and draws color-flow lines, with color flowing counterclockwise, connecting adjacent gluons. One then deforms the color-flow lines in all possible ways to form the Feynman diagrams that contribute to this partial amplitude. The Feynman diagrams that contribute to a partial amplitude are planar. This is not due to an expansion in $1/N$; the partial amplitudes are exact. They note (see their Table 1) that the number of Feynman diagrams contributing to an n -gluon partial amplitude grows as $\approx 3 \cdot 8^n$. In contrast, the number of Feynman diagrams contributing to the full amplitude grows factorially, as $\approx (2n)!$.

2017-06-09 Predrag Henn *et al.* [86] *Four-loop photon quark form factor and cusp anomalous dimension in the large- N_c limit of QCD*, [arXiv:1612.04389](https://arxiv.org/abs/1612.04389), is a thoroughly modern paper, a listing of different codes used to generate diagrams and evaluate integrals.

2017-06-16 Predrag Chang, Liu and Roberts [33] *Dressed-quark anomalous magnetic moments*

2017-06-16 Predrag Choudhury and Lahiri [36] *Anomalous chromomagnetic moment of quarks*

2017-06-16 Predrag Bermudez *et al.* [18] *Quark-gluon vertex: A perturbation theory primer and beyond*, [arXiv:1702.04437](https://arxiv.org/abs/1702.04437): The on-shell limit enables us to compute anomalous chromomagnetic moment of quarks.

[...] we present some “physically” relevant results for the on-shell limit $p^2 = k^2 = m^2$ and $q^2 = 0$. The Dirac and Pauli form factors, $F_1(q^2)$ and $F_2(q^2)$, respectively, define the Gordon decomposition of the quark current as in (1.3). The anomalous chromomagnetic moment (ACM) of quarks can be identified as $F_2(q^2)$ for $q^2 \rightarrow 0$. The Abelian version of this decomposition with $C_F = 1$ and $C_A = 0$ is the electron-photon vertex of quantum electrodynamics. The great

successes of the Dirac equation is the prediction of the magnetic moment of a charged fermion $\mu = eg/(2m)S$. The radiative corrections lead to [156]

$$\frac{e}{2m} \Rightarrow \left(1 + \frac{\alpha}{2\pi}\right) \frac{e}{2m}. \quad (1.39)$$

Note that the quark-gluon vertex differs from the electron-photon vertex already at one loop, by the contributions of an additional Feynman diagram, involving the triple-gluon vertex. In fact, apart from introducing additional color structure, this non-Abelian diagram introduces, at the one-loop level, a kinematical structure which is absent in the QED.

It is straightforward to see that for the soft gluon limit, $q^2 = 0$, the Abelian contribution for the ACM reduces to the non-Abelian counterpart of Schwinger's result, $F_2^a(0) = -\alpha/12\pi$, already derived in ref. [33]. On the other hand, the corresponding non-Abelian contribution yields a divergence [36], for a non-zero quark mass, $m \neq 0$. We find this divergence to be logarithmic. For deep infrared gluon momenta it behaves as $F_2^b(q^2 \rightarrow 0) = C_b \ln(-q^2/m^2)$. Of course, perturbation theory in QCD is not the way to explore deep infrared region. All perturbative conclusions will be taken over by non-perturbative effects, overshadowing this divergence.

2017-06-16 Predrag Brambilla *et al.* [27] *QCD and strongly coupled gauge theories: challenges and perspectives*

2017-06-16 Predrag Simonov and Tjon [163] *The Feynman–Schwinger representation in QCD*: “The proper time path integral representation is derived for Green’s functions in QCD. After an introductory analysis of perturbative properties, the total gluonic field is separated into a nonperturbative background and valence gluon part. For nonperturbative contributions the background perturbation theory is used systematically, yielding two types of expansions. As an application, we discuss the collinear singularities in the Feynman–Schwinger representation formalism.”

2017-06-16 Predrag Kälin [97] *Cyclic Mario worlds – color-decomposition for one-loop QCD*, [arXiv:1712.03539](https://arxiv.org/abs/1712.03539): “a new color decomposition for QCD amplitudes at one-loop level [...] Starting from a minimal basis of planar primitive amplitudes we write down a color decomposition that is free of linear dependencies among appearing primitive amplitudes or color factors. [...] The standard $SU(N)$ trace-based color decomposition at tree level does not take advantage of all linear dependencies of primitive amplitudes and color factors, as the sum goes over an overcomplete set of linear dependent primitive amplitudes and color factors. [...] we propose a compact color decomposition that eliminates linear dependencies of one-loop QCD amplitudes”

He uses “Melia’s basis,” “Dyck words.”

1.10 Is QED finite? A blog

1.10.1 Quenched QED/QFT

2013-11-24 Predrag As far as I can tell, terminology “of *quenched* type” was first introduced in this context by Marinari, Parisi and Rebbi [126] *Monte Carlo simulation of the massive Schwinger model*, in the context of lattice gauge theory. They write: “A first approximation to the effect of the gauge field on the fermion observables may be achieved by [...] neglecting the contributions from the fermionic vacuum polarization diagrams and, using a terminology developed in the theory of condensed matter, we shall call the expectation values thus obtained ‘quenched’”,

Parisi is reputable, so in the “quenched approximation” one neglects the fermionic vacuum polarization effects (i.e, the fermion loops) from the fermion determinant in the effective action. If it is good enough for Kinoshita, it is good enough for you.

“In the ‘quenched approximation’ the quark determinant is set equal to unity, i.e., neglecting the effect of virtual quark loops. In other words, this extreme approximation in terms of heavy quarks with a vanishing number of flavors assumes that gauge fields affect quarks while quarks have no dynamical effect on gauge fields.”

A more general usage: “In Quantum Cosmology “quenching,” amounts to quantizing a single scale factor thereby selecting a class of cosmological models, for instance, the Friedmann-Robertson-Walker space-time while neglecting the quantum fluctuations of the full metric.”

But I still do not like it - it is mostly associated with Kogut, where it means something different (as in “... treating the gauge interaction in the quenched, planar (ladder) approximation”); search for quench [here](#). Or here is what Brezin says:

“concept of quenching is well-known in the statistical mechanics of random media ; consider a system of particles, for instance, an electron gas, interacting with impurities. If these impurities are mobile, they will thermalize with the electron gas and the average physical quantities are obtained by a trace over the electron gas and the impurities degrees of freedom. However if the impurities are frozen, the ‘quenched’ case, the physical observables are obtained by calculating their value for fixed impurities and then averaging over these impurities. ”

That is how I know it - nothing about fermion loops, just dirt physics...

From my point of view, the question is whether the sum of all corrections to $(g-2)$ is a convergent series, or an asymptotic one. If one can prove the convergence for the quenched sector, I would expect each un-quenched sector (diagrams with one, two, loops) separately to be convergent, and their sum as well.

Here is something to amuse you: [on amplituhedron](#). More serious: Lance Dixon [on calculating amplitudes](#).

1971-08-01 Lautrup, Peterman, and de Rafael 1972 *Recent developments in the comparison between theory and experiments in quantum electrodynamics* [122] list the 3-loop, no-electron loop “gauge invariant subclasses” (their Fig. 4.3).

The inventor of the “gauge invariant diagram sets” concept is Benny Lautrup.

1974-01-07 Samuel 1974 *Estimates of the eighth-order corrections to the anomalous magnetic moment of the muon* [146]:

“We speculate that in making radiative corrections to a class of graphs by inserting a single photon in all possible ways, one obtains a contribution which is roughly $-\frac{\alpha}{\pi}$ times the contribution of the class. This seems to be obeyed by the known contributions.”

1.10.2 Is QED finite? A blog, continued

1950-12-22 Schwinger *On gauge invariance and vacuum polarization* [157] has 3700 citations. He writes: “ This paper is based on the elementary remark that the extraction of gauge invariant results from a formally gauge invariant theory is ensured if one employs methods of solution that involve only gauge covariant quantities. We illustrate this statement in connection with the problem of vacuum polarization by a prescribed electromagnetic field. The vacuum current of a charged Dirac field, which can be expressed in terms of the Green’s function of that field, implies an addition to the action integral of the electromagnetic field. Now these quantities can be related to the dynamical properties of a “particle” with space-time coordinates that depend upon a proper-time parameter. The proper-time equations of motion involve only electromagnetic field strengths, and provide a suitable gauge invariant basis for treating problems. Rigorous solutions of the equations of motion can be obtained for a constant field, and for a plane wave field. A renormalization of field strength and charge, applied to the modified lagrange function for constant fields, yields a finite, gauge invariant result which implies nonlinear properties for the electromagnetic field in the vacuum. [...] one can employ an expansion in powers of the potential vector. The latter automatically yields gauge invariant results, provided only that the proper-time integration is reserved to the last. This indicates that the significant aspect of the proper-time method is its isolation of divergences in integrals with respect to the proper-time parameter, which is independent of the coordinate system and of the gauge. The connection between the proper-time method and the technique of “invariant regularization” is discussed. Incidentally, the probability of actual pair creation is obtained from the imaginary part of the electromagnetic field action integral. Finally, as an application of the Green’s function for a constant field, we construct the mass operator of an electron in a weak, homogeneous external field, and derive the additional spin magnetic moment of $\alpha/2\pi$ magnetons by means of a perturbation calculation in which proper-mass plays the customary role of energy. ”

“ A proper time wave equation, in conjunction with the second order Dirac operator, has been discussed by V. A. Fock [71] *Proper time in classical and quantum*

mechanics. See also Nambu [131] (1950). ”

In Appendix B Schwinger computes the anomalous spin magnetic moment $\alpha/2\pi$ produced by second-order electromagnetic mass effects.

1977-03-03 Drell and Pagels *Anomalous magnetic moment of the electron, muon, and nucleon* [56] attempt got the sign right, but was not successful in predicting the magnitude of the sixth-order magnetic moment; $0.15 \left(\frac{\alpha}{\pi}\right)^3$ instead of $1.19 \left(\frac{\alpha}{\pi}\right)^3$.

2013-12-08 Predrag to Piotr, Wanda and Andrea (Piotr Czarski <piotr.czarski@ifj.edu.pl>, wanda.alberico@to.infn.it, andrea.prunotto@gmail.com):

I’m no fan of Feynman diagrams (my rant is [here](#)), and I’m always looking for other ways to look at perturbative expansions. So just a little email - if you have a new angle [136] on subsets of diagrams which are gauge invariant sets, I would be curious to learn how you look at that.

Just something to keep in mind :)

PS to Andrea: I realize you might rather forget this stuff (takes you a decade to write a paper?) but at least I got a ringtone out of you. The only problem is, I do not have a cell phone, so I do not know how to make it ring. At least I’m more technologically savvy than [Peter Higgs](#).

2013-12-10 Andrea Sorry for late reply (well, we’re used to longer gaps). Yes! I actually took 10 years to write this paper out of my master thesis, but I have some excuses: I did my PhD in Biochemistry (Zürich) and now I work on genetics (Lausanne). This summer my “old” professor Wanda found my work in a drawer and then contacted me, telling me that it would be a good idea to publish it.

About your request: I’m really interested in seeing if the rooted-map approach to Feynman diagrams can address the problem you’ve risen. But I have no idea what the ”subsets of diagrams which are gauge invariant sets” are. I’ve checked a bit on the web but I’m sure you can give me better indications (the works I found were too technical: I need to know the basis of the problem). Can you send me some specific link at freshman level, in particular where I can see the geometry of these subclasses of diagrams?

2013-12-11 Predrag Googling is good, but it is faster to click on [this link](#). The article defines the gauge invariant sets.

2014-02-11 M. Borinsky *Feynman graph generation and calculations in the Hopf algebra of Feynman graphs* [25] “Programs for the computation of perturbative expansions of quantum field theory amplitudes are provided. feynngen can be used to generate Feynman graphs for Yang-Mills, QED and phi^k theories. feyn-cop implements the Hopf algebra of those Feynman graphs which incorporates the renormalization procedure necessary to calculate finite results in perturbation theory of the underlying quantum field theory. ”

2016-02-08 Predrag Prunotto [135] *A Homological Approach to Feynman Diagrams in the Quantum Many-Body Theory*, and Prunotto, Alberico and Czarski [136]

2013 *Feynman Diagrams and Rooted Maps* has been submitted to the European Physical Journal A as manuscript ID EPJA-103480, seems not to have been published anywhere by 2017. They write: “ The Rooted Maps Theory, a branch of the Theory of Homology, is shown to be a powerful tool for investigating the topological properties of Feynman diagrams, related to the single particle propagator in the quantum many-body systems. The numerical correspondence between the number of this class of Feynman diagrams as a function of perturbative order and the number of rooted maps as a function of the number of edges is studied. A graphical procedure to associate Feynman diagrams and rooted maps is then stated. Finally, starting from rooted maps principles, an original definition of the genus of a Feynman diagram, which totally differs from the usual one, is given. ”

2017-03-15 Predrag Dunne and Krasnansky [57] 2006 “*Background field integration-by-parts*” and *the connection between one-loop and two-loop Heisenberg-Euler effective actions*: “ We develop integration-by-parts rules for diagrams involving massive scalar propagators in a constant background electromagnetic field, and use these to show that there is a simple diagrammatic interpretation of mass renormalization in the two-loop scalar QED Heisenberg-Euler effective action for a general constant background field. This explains why the square of a one-loop term appears in the renormalized two-loop Heisenberg-Euler effective action, and dramatically simplifies the computation of the renormalized two-loop effective action for scalar QED, and generalizes a previous result obtained for self-dual background fields. ”

2017-05-23 Predrag M. G. Schmidt and C. Schubert [148] 1994 *Multiloop calculations in the string-inspired formalism: the single spinor-loop in QED*, [arXiv:hep-th/9410100](#): They use the worldline path-integral Bern-Kosower formalism for to calculate the sum of all diagrams with one spinor loop and fixed numbers of external and internal photons. Of interest: in this formalism the three 2-loop photon polarization graphs, see figure 1.7, are a single integral, easier to evaluate than any of the three Feynman graphs. They also note an unexplained cancellation not only of poles, but also of “transcendentals.” A knot-theoretic explanation for the rationality of the quenched QED beta function is given in ref. [29].

2017-05-23 Predrag Nieuwenhuis and Tjon [132] 1996 *Nonperturbative study of generalized ladder graphs in a $\phi^2\chi$ theory*, [arXiv:hep-ph/9606403](#):

2017-05-23 Predrag Christian Schubert [152] 2001 *Perturbative quantum field theory in the string-inspired formalism*, [arXiv:hep-th/0101036](#):

The Feynman rules for (Euclidean) spinor QED in the second order formalism (see Morgan [130] 1995, Strassler [169] 1992, and references therein) are, up to statistics and degrees of freedom, the ones for scalar QED with the addition of a third vertex. The third vertex involves $\sigma^{\mu\nu} = \frac{1}{2}[\gamma^\mu, \gamma^\nu]$ and corresponds to the $\psi^\mu F_{\mu\nu} \psi^\nu$ – term in the worldline Lagrangian L_{spin} . For the details and for the non-abelian case see Morgan [130]. There also an algorithm is given, based

on the Gordon identity, which transforms the sum of Feynman (momentum) integrals resulting from the first order rules into the ones generated by the second order rules.

2017-06-12 Predrag Morgan [130] *Second order fermions in gauge theories*, [arXiv:hep-ph/9502230](#) seems to be the same discussion that I use in my QFT course to define the electron magnetic moment via $\sigma^{\mu\nu}$ (see lectures 25 and 26 [here](#))

2017-06-12 Predrag Gies, Sanchez-Guillen, Vázquez [84] *Quantum effective actions from nonperturbative worldline dynamics*, [arXiv:hep-th/0505275](#): “ We demonstrate the feasibility of a nonperturbative analysis of quantum field theory in the worldline formalism with the help of an efficient numerical algorithm. In particular, we compute the effective action for a super-renormalizable field theory with cubic scalar interaction in four dimensions in quenched approximation (small- N expansion) to all orders in the coupling. We observe that nonperturbative effects exert a strong influence on the infrared behavior, rendering the massless limit well defined in contrast to the perturbative expectation. ”

2017-06-12 Predrag Gies and Hämmerling [82] *Geometry of spin-field coupling on the worldline*, [arXiv:hep-th/0505072](#):

2017-03-15 Predrag Huet, McKeon, and Schubert [88] 2010 *Euler-Heisenberg Lagrangians and asymptotic analysis in 1+1 QED. Part I: Two-loop* (no GaTech online access, [arXiv:1010.5315](#)): “ We continue an effort to obtain information on the QED perturbation series at high loop orders, and particularly on the issue of large cancellations inside gauge invariant classes of graphs, using the example of the 1-loop N -photon amplitudes in the limit of large photon numbers and low photon energies. The high-order information on these amplitudes can be obtained from a nonperturbative formula, due to Affleck *et al.* [1], for the imaginary part of the QED effective Lagrangian in a constant field. The procedure uses Borel analysis and leads, under some plausible assumptions, to a number of nontrivial predictions already at the three-loop level. Their direct verification would require a calculation of this ‘Euler-Heisenberg Lagrangian’ at three-loops, which seems presently out of reach (though see Huet, de Trautenberg, and Schubert [90] below). Motivated by previous work by Dunne and Krasnansky [57] on Euler-Heisenberg Lagrangians in various dimensions, in the present work we initiate a new line of attack on this problem by deriving and proving the analogous predictions in the simpler setting of 1+1 dimensional QED. In the first part of this series, we obtain a generalization of the formula of Affleck *et al.* [1] to this case, and show that, for both scalar and spinor QED, it correctly predicts the leading asymptotic behaviour of the weak field expansion coefficients of the two loop Euler-Heisenberg Lagrangians.

“The present work continues an effort [58–60, 127] to study the multiloop behaviour of the QED N -photon amplitudes using the QED effective Lagrangian, and in particular to prove or disprove Cvitanović’s conjecture for these amplitudes.”

2017-05-24 Predrag Bastianelli, Huet, Schubert, Thakur and Weber [17] 2014 *Integral representations combining ladders and crossed-ladders* write:

This property is particularly interesting in view of the fact that it is just this type of summation which in QED often leads to extensive cancellations, and to final results which are substantially simpler than intermediate ones (see, e.g., ref. [29, 41]). More recently, similar cancellations have been found also for graviton amplitudes (see, e.g., ref. [13]). Although this property of the worldline formalism is well-known, and has been occasionally exploited [15, 141, 142, 148] (see also ref. [76]) a systematic study of its implications is presently still lacking.

The first classes of Green's functions is the x -space propagator for one scalar interacting with the second one through the exchange of N given momenta.

This object, to be called " N -propagator", is given by a set of $N!$ simple tree-level graphs, is in the worldline formalism combined into a single integral.

The second class are the similarly looking x -space $N + 2$ - point functions defined by a line connecting the points x and y and N further points z_1, \dots, z_N connecting to this line in an arbitrary order.

An advantage of the worldline representation over the usual Feynman parameterization is the automatic inclusion of all possible ways of crossing the "rungs" of the ladders. They obtain such representations in explicit form both in x -space and in momentum space.

The inclusion of the crossed ladder graphs is essential for the consistency of the one-body limit where one of the constituents becomes infinitely heavy, and for maintaining gauge invariance.

They concentrate on the case of infinite N , *i.e.*, the sum over *all* ladder *and* crossed ladder graphs.

As their main application, they consider the case of two massive scalars interacting through the exchange of a massless scalar, obtain an the case of a massless exchanged particle (along the "rungs" of the ladders).

Applying asymptotic estimates and a saddle-point approximation to the N -rung ladder plus crossed ladder diagrams, they derive a semi-analytic approximation formula for the lowest bound state mass in this model.

They use the worldline formalism to derive integral representations for the N -propagators and the N -ladders - in scalar field theory, and give a compact expression combining the $N!$ Feynman diagrams contributing to the amplitude. They give these representations in both x and (off-shell) momentum space. Being off-shell, can be used as building blocks for more complex amplitudes. They derive a compact expression for the sum of all ladder graphs with N rungs, including all possible crossings of the rungs.

Nieuwenhuis and Tjon [132] 1996 have numerically evaluated the path integrals of the worldline representation for the same scalar model field theory, thus including all ladder *and* crossed ladder graphs.

2017-05-23 Predrag Huet, de Traubenberg, and Schubert [90] 2017 *Multiloop Euler-Heisenberg Lagrangians, Schwinger pair creation, and the photon S-matrix*: “Schwinger pair creation in a constant electric field, may possibly provide a window to high loop orders; simple non-perturbative closed-form expressions have been conjectured for the pair creation rate in the weak field limit, for scalar QED in 1982 by Affleck, Alvarez, and Manton [1], and for spinor QED by Lebedev and Ritus [123] in 1984. Using Borel analysis, these can be used to obtain non-perturbative information on the on-shell renormalized N-photon amplitudes at large N and low energy.”

“there is something quite implausible about it: a summation to all loop orders has produced the perfectly analytic factor $e^{\alpha\pi}$! This is certainly contrary to standard QED wisdom”

Preliminary results of a calculation of the three-loop Euler–Heisenberg Lagrangian in two dimensions indicate that the exponentiation conjecture by Affleck *et al.* and Lebedev/Ritus probably fails in $D = 2$.

Dunne and Schubert conjectured in 2005 that the QED N –photon amplitudes in the quenched (one electron loop) approximation are convergent in perturbation theory [60]. In this article they say; “Later they learned that Cvitanović in 1977 had already made the analogous conjecture for $(g - 2)$ [41].”

2017-06-14 Predrag Academician Ritus [139] writes: “The requirement $e^2/\hbar c = 1$ leads to unique values of the point-like charge and its fine structure constant, $e_0 = \pm\sqrt{\hbar c}$, $\alpha_0 = 1/4\pi$. Arguments are adduced in favor of the conclusion that this value of the fine structure constant is the bare, nonrenormalized value.”

In 1951 Ritus was assigned to what was known at the time as the First Main Directorate of the USSR Council of Ministers, later rechristened the Ministry of Medium Machine Building (Sredmash) – a powerful state body placed above any other in the name of implementing the Soviet Government sponsored program of thermonuclear weapons design. [...] The legend of his infallibility when conducting complicated and cumbersome computations, just started to take root; its protagonist did nothing that would sully this reputation, neither then nor later. Science is unaware of any errors ever made by Ritus! (from ref. [23].)

2017-05-23 Predrag Das, Frenkel and Schubert [53] 2013 *Infrared divergences, mass shell singularities and gauge dependence of the dynamical fermion mass*; [arXiv:1212.2057](https://arxiv.org/abs/1212.2057):

2017-05-23 Predrag Ahmadinia, Bashir and Schubert [3] 2016 *Multiphoton amplitudes and generalized Landau-Khalatnikov-Fradkin transformation in scalar QED*, [arXiv:1511.05087](https://arxiv.org/abs/1511.05087):

$D_{\mu\nu}$ is the x -space photon propagator. In D dimensions and arbitrary covariant gauge

$$D_{\mu\nu}(x) = \frac{1}{4\pi^{\frac{D}{2}}} \left\{ \frac{1+\xi}{2} \Gamma\left(\frac{D}{2}-1\right) \frac{\delta_{\mu\nu}}{(x^2)^{\frac{D}{2}-1}} + (1-\xi) \Gamma\left(\frac{D}{2}\right) \frac{x_\mu x_\nu}{(x^2)^{\frac{D}{2}}} \right\}. \quad (1.40)$$

Calculation methods:

1. The analytic or “string-inspired” approach, based on the use of worldline Green’s functions: all path integrals are brought into Gaussian form; this requires some expansion and truncation. They are then calculated by Gaussian integration.
2. The semi-classical approximation, based on a stationary trajectory (“world-line instanton”).

We will focus on the closed-loop case in the following, since it turns out to be simpler than the propagator one. Nevertheless, it should be emphasized that everything that we will do in the following for the effective action can also be done for the propagator.

Some reasonable gymnastics leads to the “Bern-Kosower master formula” [19, 20, 169]

2017-05-23 Predrag Strassler [169] *Field theory without Feynman diagrams: One-loop effective actions*, [arXiv:hep-ph/9205205](#);

2017-05-23 Predrag Ahmad *et al.* [2] 2017 *Master formulas for the dressed scalar propagator in a constant field*, [arXiv:1612.02944](#)

2007-01-31 Kurusch Ebrahimi-Fard Here are the links I mentioned:

Anatomy of a gauge theory by Dirk Kreimer, [arXiv:hep-th/0509135](#)

Renormalization of gauge fields: A Hopf algebra approach by Walter D. van Suijlekom, [arXiv:hep-th/0610137](#)

The Hopf algebra of Feynman graphs in QED by Walter D. van Suijlekom, [arXiv:hep-th/0602126](#)

This is Jean-Yves Thibon’s [web-page](#), a very good combinatorialist!

2016-11-15 Kevin Hartnett [Strange Numbers Found in Particle Collisions](#)

2017-05-23 Predrag Should I talk to [Spencer Bloch](#)?

2017-05-23 Predrag Broadhurst, Delbourgo and Kreimer [29] 1996 *Unknotting the polarized vacuum of quenched QED* has lots of magic leading to cancelations of “transcendentals.” They say: “Complete cancellation of transcendentals from the beta function, at every order, is to be expected only in quenched QED and quenched SED, where subdivergences cancel between bare diagrams.”

Online collection of papers on [this topic](#).

2013-10-23 Warren D. Smith D. J. Broadhurst and D. Kreimer: *Association of multiple zeta values with positive knots via Feynman diagrams up to 9 loops*, Physics Letters B 393 (1997) 403-412, [arXiv:hep-th/9609128](#).

Furthermore, the number of different kinds of knots with N crossings, $\text{KnotCount}(N)$, is known asymptotically to be bounded between two simple-exponentials,

$$A^N < \text{KnotCount}(N) < B^N$$

where $B \leq 13.5$ according to

D.J.A. Welsh, *On the number of knots and links, Sets, graphs and numbers* (Budapest, 1991), 713–718, Colloq. Math. Soc. Janos Bolyai, 60, North-Holland, Amsterdam, 1992,

while $A \geq 2.68$ according to

C.Ernst & D.W.Summers *The growth of the number of prime knots*, Proc Cambridge Philo Soc 102 (1987) 303-315.

So, that's the funny thing. My attempt to further-destroy Cvitanović, just led to an estimate involving simple exponential growth and NOT superexponential (e.g. factorial style). This is right on the boundary for convergence questions, i.e.

$$\sum A_N x^N$$

has a finite, nonzero radius of convergence if $|A_N|$ grows exponentially. So maybe there remains some hope for some form of Cvitanović conjecture.

The Welsh upper bound also works for links. Which means: if you believe this could rescue Cvitanović's quenched-QED convergence conjecture, that would also presumably mean you believe full unquenched QED series have finite nonzero radius of convergence.

2013-11-25 David Broadhurst <David.Broadhurst@open.ac.uk>

Dirk and kind of gave up when it turned out that a pair of counterterms at 7 loops, unidentified in 1996, have weight 11, whereas our intuition about the knots 10_139 and 10_154 has suggested weight 10.

Maybe there is some sort of connection, but I know not what.

2017-05-23 Predrag Dirk Kreimer and **Karen Yeats** [115] 2008 *Recursion and growth estimates in renormalizable quantum field theory*

Our method is very different in spirit from the constructive approach or the functional integral approach. It relies on a Hopf algebraic decomposition of terms in the perturbative expansion into primitive constituents, not unlike the decomposition of a ζ function into Euler factors.

Our construction of a basis of primitives with a given Mellin transform resolves overlapping divergences, thanks to the Hochschild cohomology of the relevant Hopf algebras [113].

We next assume there to be $p(k)$ primitives at k loops where p is a polynomial.

Yeats [181] 2017 *A Combinatorial Perspective on Quantum Field Theory*. I have put a copy [here](#).

2016-08-20 Predrag Kießler [106] *Hopf-algebraic renormalization of QED in the linear covariant gauge*: “The possibility of a finite electron self-energy by fixing a generalized linear covariant gauge is discussed. An analysis of subdivergences leads to the conclusion that such a gauge only exists in quenched QED.”

2017-06-16 Henry Kißler The term “finite electron self-energy” does not refer to the convergence of the perturbation series, but to an order-by-order cancellation of divergences. The idea was to gauge away all divergences in the self-energy order-by-order as used by Broadhurst [28] in “Four-loop Dyson-Schwinger-Johnson anatomy” for quenched QED.

2017-06-16 Predrag Broadhurst [28] *Four-loop Dyson-Schwinger-Johnson anatomy*: “Dyson–Schwinger equations are used to evaluate the 4-loop anomalous dimensions of quenched QED in terms of finite, scheme-independent, 3-loop integrals. The 4-loop beta function has 24 unambiguous terms. The rational, $\zeta(3)$ and $\zeta(5)$ parts of the other 22 miraculously sum to zero. Vertex anomalous dimensions have 40 terms, with no dramatic cancellations. Our methods come from work by the late Kenneth Johnson, done more than 30 years ago. They are entirely free of the subtractions and infrared rearrangements of later methods.”

2016-08-20 Predrag Kißler and Kreimer [107] 2016 *Diagrammatic cancellations and the gauge dependence of QED*, [arXiv:1607.05729](https://arxiv.org/abs/1607.05729): “The perturbative expansion given in terms of Feynman graphs might be rearranged in terms of meta graphs or subsectors with a maximum number of cancellations implemented.”

“The summation over all insertions requires considering *connected* rather than one-particle irreducible Feynman graphs.”

In his (unpublished) thesis Henry defines the above “self-energy set” as an “equivalence class” obtained by all insertions of an external electron-photon vertex into the external electron line propagator of a self-energy graph. On mass-shell the summation over this class is gauge invariant by the Ward-Takahashi identity [172, 180].

This gets more interesting when he considers insertion of internal photon propagator “gaugeons” (a Lautrup name?). Then, for example (his eq. (2.75)) the sum of all connected quenched 2-loop graphs is equivalent to the one-loop self energy. Perhaps iteration of such insertions into Schwinger 1-loop anomaly might offer a proof of invariance of gauge sets...

“The four-photon interaction is called light-by-light scattering and supposed to be finite by renormalizability –in other words there is no interaction vertex in Quantum Electrodynamics that could serve to absorb the divergences of the four-photon type. Unfortunately, the author is not aware of a general argument beyond the first-loop order that proves this statement. In [arXiv:1406.1618](https://arxiv.org/abs/1406.1618) and [arXiv:1703.01094](https://arxiv.org/abs/1703.01094), a basis of Lorentz tensors was constructed for the class four-photon graphs; it consists of 138 Lorentz tensors and the challenge is to show that each coefficient becomes finite when all graphs at a certain loop order are considered.”

He shows that the sum over all one-particle irreducible four-photon graphs is transversal (it vanishes whenever one of its external legs is contracted with the associated momentum). This global property restricts the possible occurrence of divergences to 43 Lorentz tensors (see [arXiv:1505.06336](https://arxiv.org/abs/1505.06336) for an alternative basis

of Lorentz tensors which includes the anti-symmetry tensor), whose coefficients remain to be proven finite.

They discuss how the QED tree-level cancellation identity implies cancellation between Feynman graphs of different topologies and determines the gauge dependence. They parameterize the momentum part of Landau gauge, then start by keeping only a linear term in graphs (i.e., insert only one Landau propagator, rest Feynman). They focus on the electron propagator in the massless limit of Quantum Electrodynamics. Not sure it is useful to us...

2017-06-16 Henry Your definition of gauge sets differs from the one we use in *Diagr. cancellations and the gauge dependence* [107]. The anomalous mag. moment is gauge independent due to the on-shell electrons, so studying the gauge parameter terms is not important, but I find it interesting to compare both definitions of gauge invariant sets, maybe one can improve the other.

Read also Kreimer [114] 2000 *Knots and Feynman diagrams*.

2017-06-16 Henry Kibler Here an alternative idea to approach the finiteness conjecture: There is a bound for the value of a scalar Feynman diagram due to Eric Panzer, which he called the *Hepp bound*. This bound is derived using the 1974 Cvitanović and Kinoshita [48] Feynman-parametric representation. A natural question is: does this bound generalize to a sum of Feynman graphs when the sum goes over something as your gauge sets. Of course, the gamma matrices in the numerator make things more complicated, but imposing on-shell conditions and choosing an appropriate gauge might simplify this task. Unfortunately, there is little in the literature about the Hepp bound [151]; the only thing I am aware of is a short section in the [thesis](#) of Iain Crump [37].

2017-06-16 Predrag I do not see how this would work: The Hepp invariant $H(G)$ is defined for a given graph G . Suppose we use instead of a Feynman diagram the $km'm$ multi-photons gauge set diagram \tilde{G} (for examples, see figure 1.4 and figure 1.5). In worldline formalism one has a proper time parametrization intermingled with Feynman-parametric bits. Any clue how the weights in the Hepp invariant $H(\tilde{G})$ would be defined? Just for a scalar theory, let us forget QED for the time being, along the lines of **2017-05-24 Predrag** entry above, on Bastianelli *et al.* [17]?

2017-06-19 Predrag Iain Crump [37] [thesis](#) *Graph Invariants with Connections to the Feynman Period in ϕ^4 Theory* (he follows Yeats [181]):

The *Feynman period* is a simplified version of the Feynman integral. The period is of special interest, as it maintains much of the important number theoretic information from the Feynman integral. It is also of structural interest, as it is known to be preserved by a number of graph theoretic operations.

2017-05-23 Predrag Badger, Bjerrum-Bohr and Vanhove [13] 2009 *Simplicity in the structure of QED and gravity amplitudes*.

2017-05-23 Predrag Rosenfelder and Schreiber [141, 142] 1996, [arXiv:nucl-th/9504002](#), [arXiv:nucl-th/9504005](#):

2017-06-02 Predrag Rosenfelder and Schreiber [143] 2004 *An Abraham-Lorentz-like equation for the electron from the worldline variational approach to QED*:

They discuss the of a spin-1/2 electron dressed by an arbitrary number of photons in the quenched approximation to QED. The approach is patterned after Feynman’s celebrated variational treatment of the polaron problem [69], which was first applied by Mano, *Progr. Theor. Phys.* 14, 435 (1955) [8] to a relativistic scalar field theory and rediscovered and expanded by them in a series of papers [141, 142]. Its main features are the description of relativistic particles by worldlines [152] parametrized by the proper time, an exact functional integration over the photons and a variational approximation of the resulting effective action by a retarded quadratic trial action. In recent work we have extended this approach to more realistic theories, in particular to quenched QED [4] (the divergence structure and renormalization, a compact expression for the anomalous mass dimension of the electron). Here they calculate the finite contributions.

The variational formulation of worldline QED leads to an equation which is similar to Abraham, Lorentz and Dirac description of the electron and its self-interaction with the radiation field. The approach contains (almost) all the ingredients of the relativistic field theory of electrons and photons, in particular its divergence structure. This has been demonstrated by deriving an approximate nonperturbative expression for the anomalous mass dimension of the electron.

2017-05-23 Predrag K. Barro-Bergflödt, R. Rosenfelder and M. Stingl [15] 2006 *Variational worldline approximation for the relativistic two-body bound state in a scalar model*, [arXiv:hep-ph/0601220](https://arxiv.org/abs/hep-ph/0601220).

2017-05-23 Predrag Fried and Gabellini [75] 2009 *Analytic, nonperturbative, almost exact QED: The two-point functions*. The remarkable (but speculative) result of this paper is that in a convenient gauge, the (unphysical) electron propagator renormalization is a multiplicative, non-perturbative *finite* factor bounded between 0 and 1:

$$Z_2 = \exp \left[-2\gamma \left(\left(\frac{\pi}{2} \right)^2 + \ln^2 \left(\frac{\Lambda^2}{\mu^2} \right) \right) \right], \quad (1.41)$$

where $\gamma = e^2/4\pi^2$ is the fine structure constant (referred to by the vulgar multitudes as α), $\mu \rightarrow 0$ is the infrared cutoff, and $\Lambda \rightarrow \infty$ is the UV cutoff. One would still need to compute the vertex renormalization Z_1 to get a gauge and renormalization method invariant result. Fried seem to only cite Schwinger, Fradkin and himself, so the similarity of this to 1982 Affleck, Alvarez, and Manton [1] nonperturbative result $e^{\alpha\pi}$ is not remarked upon.

Fried and Gabellini [76] 2012 *On the Summation of Feynman Graphs*, [arXiv:1004.2202](https://arxiv.org/abs/1004.2202).

Fried and Gabellini [77] 2013 *QED vacuum loops and vacuum energy*

2017-03-15 Predrag I have tried reading Fried [74] 2014 *Modern Functional Quantum Field Theory: Summing Feynman Graphs*: “a simple, analytic, functional approach to non-perturbative QFT, using a functional representation of Fradkin

to explicitly calculate relevant portions of the Schwinger Generating Functional (GF). In QED, this corresponds to

summing all Feynman graphs representing virtual photon exchange

between charged particles. It is then possible to see, analytically, the cancellation of an infinite number of perturbative, UV logarithmic divergences, leading to an approximate but most reasonable statement of finite charge renormalization. A similar treatment of QCD, with the addition of a long-overlooked but simple rearrangement of the Schwinger GF which displays Manifest Gauge Invariance, is then able to produce a simple, analytic derivation of quark-binding potentials without any approximation of infinite quark masses. A crucial improvement of previous QCD theory takes into account the experimental fact that asymptotic quarks are always found in bound state.”

This book can be read online via GaTech library link [here](#) or [here](#).

Even though I am a grandchild of Schwinger (via Tung Mow Yan), and have written/drawn a book where Schwinger’s functional formalism is explained to everywoman, I still find the functional formalism of Schwinger and Fradkin [72] hard to follow. I believe the results are essentially the same as wordline formalism developed by Schubert *et al.*

2017-06-15 Predrag I’ve now reread much of the relevant literature known to me. There might be much more - people who do things related to my 1977 paper [41] never alert me to their papers, presumably because the reports of my death have been greatly exaggerated.

2017-06-16 Predrag Jia and Pennington [93] *How gauge covariance of the fermion and boson propagators in QED constrain the effective fermion-boson vertex*

Jia and Pennington [94] *Gauge covariance of the fermion Schwinger–Dyson equation in QED*

Jia and Pennington [95] *Landau-Khalatnikov-Fradkin transformation for the fermion propagator in QED in arbitrary dimensions*

2017-06-16 Christian Melnikov, Vainshtein and Voloshin [128] *Remarks on the effect of bound states and threshold in $g=2$* : “The appearance of positronium poles in a photon propagator in QED formally requires a summation of an infinite series of terms in perturbation theory. [...] we show how these nonperturbative contributions disappear, using the case of the electron anomalous magnetic moment as an example. [...] it never happens that a summation of infinite classes of Feynman diagrams enhanced at any threshold generates additional effects beyond perturbation theory. The misunderstanding of this fact appears to be quite common.”

The same conclusion is reached by Eides [63] *Recent ideas on the calculation of lepton anomalous magnetic moments* and Fael and Passera [66] *Positronium contribution to the electron $g - 2$* .

2017-06-16 Predrag Herzog and Ruijl [87] *The R*-operation for Feynman graphs with generic numerators*: “The R*-operation by Chetyrkin, Tkachov, and Smirnov is a generalisation of the BPHZ R-operation, which subtracts both ultraviolet and infrared divergences of euclidean Feynman graphs with non-exceptional external momenta. It can be used to compute the divergent parts of such Feynman graphs from products of simpler Feynman graphs of lower loops. In this paper we extend the R*-operation to Feynman graphs with arbitrary numerators, including tensors. We also provide a novel way of defining infrared counterterms which closely resembles the definition of its ultraviolet counterpart. We further express both infrared and ultraviolet counterterms in terms of scaleless vacuum graphs with a logarithmic degree of divergence. By exploiting symmetries, integrand and integral relations, which the counterterms of scaleless vacuum graphs satisfy, we can vastly reduce their number and complexity.”

Ruijl, Ueda, Vermaseren and Vogt [145] *Four-loop QCD propagators and vertices with one vanishing external momentum*: “ We have computed the self-energies and a set of three-particle vertex functions for massless QCD at the four-loop level. The vertex functions are evaluated at points where one of the momenta vanishes. Analytical results are obtained for a generic gauge group and with the full gauge dependence, which was made possible by extensive use of the Forcer program for massless four-loop propagator integrals. The bare results in dimensional regularization are provided in terms of master integrals and rational coefficients; the latter are exact in any space-time dimension.”

Chetyrkin and Tkachov [35] *Infrared R-operation and ultraviolet counterterms in the MS-scheme*, together with We Chetyrkin and Smirnov [34] *R*-Operation corrected*

Smirnov and Chetyrkin [165] *R* operation in the minimal subtraction scheme*

Johnson and Zumino [96] *Gauge dependence of the wave-function renormalization constant in Quantum Electrodynamics*: “[...] point out the existence of an exact and simple relation between the electron Green’s function renormalization constants in the general class of manifestly covariant gauges. ”

Korthals Altes and De Rafael [111] 1976 *Infrared structure of non-abelian gauge theories: An instructive calculation*

Korthals Altes and De Rafael [112] 1977 *Infrared structure of non-abelian gauge theories: Comments on perturbation theory calculations*

Frenkel *et al.* [73] 1976 *Infra-red behaviour in non-abelian gauge theories*

2017-06-27 Predrag Søndergaard, Palla, Vattay, and Voros [167] *Asymptotics of high order noise corrections*: “We consider an evolution operator for a discrete Langevin equation with a strongly hyperbolic classical dynamics and noise with finite moments. Using a perturbative expansion of the evolution operator we calculate high order corrections to its trace in the case of a quartic map and Gaussian noise. The asymptotic behaviour is investigated and is found to be independent up to a multiplicative constant of the distribution of noise.”

2017-06-16 Predrag Read also Simonov and Tjon 1996 article (PC: cannot find any such) which applies the the so-called Fock–Feynman–Schwinger representation (FSR), based on the Fock–Schwinger [157] proper time and Feynman path integral formalism, to QED.

Simonov [161] *Relativistic path integral and relativistic Hamiltonians in QCD and QED* writes: “The proper-time 4D path integral is used as a starting point to derive the new explicit parametric form of the quark-antiquark Green’s function in gluonic and QED fields entering as a common Wilson loop. The subsequent vacuum averaging of the latter allows us to derive the instantaneous Hamiltonian. The explicit form and solutions are given in the case of the $q\bar{q}$ mesons in magnetic field.”

Simonov [162] *QED spectra in the path integral formalism* writes: “Relativistic Hamiltonians, derived from the path integrals, are known to provide a simple and useful formalism for hadron spectroscopy in QCD. The accuracy of this approach is tested using the QED systems, and the calculated spectrum is shown to reproduce exactly that of the Dirac hydrogen atom. [...] The calculated positronium spectrum, including spin-dependent terms, coincides with the standard QED perturbation theory to the considered order $O(\alpha^4)$.”

2017-08-03 Predrag Schwartz [155] **Chapter33** *Effective actions and Schwinger proper time* looks very pedagogical.

2017-08-04 Predrag Mielniczuk *et al.* [129] *The anomalous magnetic moment of a photon propagating in a magnetic field*

Valluri, Jentschura and Lamm [176] *The study of the Heisenberg-Euler Lagrangian and some of its applications*

2017-08-18 Predrag Antonov [6] *World-line formalism: Non-perturbative applications*

2017-09-17 Predrag Schäfer and I. Huet and H. Gies [147] *Worldline numerics for energy-momentum tensors in Casimir geometries*: “We develop the worldline formalism for computations of composite operators such as the fluctuation induced energy-momentum tensor. As an example, we use a fluctuating real scalar field subject to Dirichlet boundary conditions. The resulting worldline representation can be evaluated by worldline Monte-Carlo methods in continuous space-time. The method generalizes straightforwardly to arbitrary Casimir geometries and general background potentials.”

2017-09-17 Predrag A closer reading of sect. 6. *Worldline formalism* of Gelis and N. Tanji [80] *Schwinger mechanism revisited* might be helpful - it goes to the barycenter to reexpress the integral as an average over Wilson loops.

In sect. 6.4. *Lattice worldline formalism* they describe a formulation of the worldline formalism the (Euclidean) space–time discretized on a cubic lattice [113–116].

Schmidt and Stamatescu [149, 150] *Matter determinants in background fields using random walk world line loops on the lattice*, using Schwinger's proper time formalism, pointed out that the fermion and boson determinant on the lattice can be viewed as a gas of closed loops which can be simulated numerically via a random walk.

Seiler and Stamatescu [160] *A note on the loop formula for the fermionic determinant*: “ A formula expressing the fermionic determinant as an infinite product of smaller determinants is derived and discussed. These smaller determinants are of a fixed size, independent of the size of the lattice and are indexed by loops of increasing length.

We will discuss in more detail the derivation of the formula as well as its limitations and possible misunderstandings in using it, since one might think that the zeroes of the determinant are given by the zeroes of the factors of the product. In fact this is not the case, and the formula should instead be understood as a systematic approximation.

In particular, for a finite lattice this formula expresses the determinant which is a polynomial of finite order in the hopping parameter as an infinite product. Obviously this can make sense only where the infinite product converges, which is equivalent to convergence of the expansion of its logarithm; this convergence will break down at the latest at the first zero encountered, either on the left-hand side (lhs) or in one of the factors on the right-hand side (rhs). ”

Turns out this is the same discussion as my discussion of the Euler product representation of dynamical ζ functions in ChaosBook.org.

Fry [78] *Nonperturbative quantization of the electroweak model's electrodynamic sector*: “ Consider the Euclidean functional integral representation of any physical process in the electroweak model. Integrating out the fermion degrees of freedom introduces 24 fermion determinants. Suppose the functional integral over the Maxwell field is attempted first. This paper is concerned with the large amplitude behavior of the Maxwell effective measure. We examine the large amplitude variation of a single QED fermion determinant. To facilitate this the Schwinger proper time representation of this determinant is decomposed into a sum of three terms. The advantage of this is that the separate terms can be non-perturbatively estimated for a measurable class of large amplitude random fields in four dimensions. It is found that the QED fermion determinant grows too fast in the absence of background potentials. Including zero mode supporting background potentials can result in a decaying exponential growth of the fermion determinant. This is prima facie evidence that Maxwellian zero modes are necessary for the nonperturbative quantization of QED ”

Gies and Langfeld [83] *Loops and loop clouds — a numerical approach to the worldline formalism in QED* point out that the fermion (and boson) determinant on the lattice can be viewed as a gas of closed loops which can be simulated numerically via a random walk: “ A numerical technique for calculating effective actions of electromagnetic backgrounds is proposed, which is based on the string-inspired worldline formalism. As examples, we consider scalar electro-

dynamics in three and four dimensions to one-loop order. Beyond the constant-magnetic-field case, we analyze a step-function-like magnetic field exhibiting a nonlocal and nonperturbative phenomenon: “magnetic-field diffusion”.

Epelbaum, Gelis and Wu [65] *From lattice Quantum Electrodynamics to the distribution of the algebraic areas enclosed by random walks on \mathbb{Z}^2* .

Epelbaum, Gelis and Wu [64] *Lattice worldline representation of correlators in a background field*: “We use a discrete worldline representation in order to study the continuum limit of the one-loop expectation value of dimension two and four local operators in a background field. We illustrate this technique in the case of a scalar field coupled to a non-Abelian background gauge field. The first two coefficients of the expansion in powers of the lattice spacing can be expressed as sums over random walks on a d-dimensional cubic lattice. Using combinatorial identities for the distribution of the areas of closed random walks on a lattice, these coefficients can be turned into simple integrals. Our results are valid for an anisotropic lattice, with arbitrary lattice spacings in each direction.”

Laufer and Orland [120] *Metric of Yang-Mills orbit space on the lattice*: “We find coordinates, the metric tensor, the inverse metric tensor and the Laplace-Beltrami operator for the orbit space of Hamiltonian SU(2) gauge theory on a finite, rectangular lattice, with open boundary conditions. This is done using a complete axial gauge fixing.”

Vilela Mendes [177] *A consistent measure for lattice Yang–Mills* (no GaTech access to the journal): “The construction of a consistent measure for Yang–Mills is a precondition for an accurate formulation of nonperturbative approaches to QCD, both analytical and numerical. Using projective limits as subsets of Cartesian products of homomorphisms from a lattice to the structure group, a consistent interaction measure and an infinite-dimensional calculus have been constructed for a theory of non-Abelian generalized connections on a hypercubic lattice. Here, after reviewing and clarifying past work, new results are obtained for the mass gap when the structure group is compact.”

2017-09-17 Predrag a letter

to: Erhard Seiler <ehs@mpp.mpg.de>,
Ion-Olimpiu Stamatescu <I.O.Stamatescu@thphys.uni-heidelberg.de>

Dear Erhard and Ion-Olimpiu

I have read your paper (well, not read it sufficiently deeply)

```
@Article{SeiStal6,
  author = {E. Seiler and Stamatescu, I. O.},
  title = {
A note on the loop formula for the fermionic determinant},
  journal = {J. Phys. A},
  year = {2016},
  volume = {49},
```

```
pages    = {335401},  
doi      = {10.1088/1751-8113/49/33/335401},  
}
```

with great interest.

2017-09-19 Schubert With g-2 we have not made a lot of headway yet, either, but my students Cesar and Misha have started chipping away at it, and with James we are making some progress in using parity and integration by parts to simplify the integrands.

2017-09-19 Schubert Arkani-Hamed, Huang and Huang [12] *Scattering amplitudes for all masses and spins*, [arXiv:1709.04891](https://arxiv.org/abs/1709.04891) contains an elegant calculation of the one-loop g-2 which shares with our approach the property that it lumps together the irreducible and the reducible diagrams.

2017-09-19 Predrag Arkani-Hamed, N. and Huang, T.-C. and Huang [12] *Scattering amplitudes for all masses and spins*: “ We introduce a formalism for describing four-dimensional scattering amplitudes for particles of any mass and spin. This naturally extends the familiar spinor-helicity formalism for massless particles to one where these variables carry an extra SU(2) little group index for massive particles, with the amplitudes for spin S particles transforming as symmetric rank 2S tensors. We systematically characterise all possible three particle amplitudes compatible with Poincare symmetry. [...]”

We illustrate a number of applications of the formalism at one-loop, giving few-line computations of the electron (g-2) as well as the beta function and rational terms in QCD. “Off-shell” observables like correlation functions and form-factors can be thought of as scattering amplitudes with external “probe” particles of general mass and spin, so all these objects—amplitudes, form factors and correlators, can be studied from a common on-shell perspective. ”

2017-12-17 Christian Schubert We have not forgotten about g-2 here, my student Misha is working on the LBL diagram, and we have also made progress in decomposing the pure fermion-line contributions, the ones that interest you most, into spin and orbit parts. My big hope is that not all will equally contribute for large N .

Last week James and I were in Bologna where we had an opportunity to talk to Remiddi. He seems not terribly interested in the convergence issue, but in any case he said he would find nothing odd if that happened to be true, since the issue of cancellations is so little understood.

Laporta has obtained a job at Padua University. More than well-deserved.

2018-01-08 Christian Schubert We are planning to have a workshop devoted to world-line methods in Germany at the Mainz Center of Theoretical Physics sometime in 2019, most likely in February. Would you be interested in principle to attend?

2018-06-02 Predrag Jegerlehner [91] *The Anomalous Magnetic Moment of the Muon* is a very detailed review of everything known about anomalous magnetic moments.

Todorov lecture [174] *From Euler's play with infinite series to the anomalous magnetic moment* is a fun overview of the history of the $(g-2)$ calculations.

2018-06-02 Predrag Aoyama and T. Kinoshita and M. Nio [11] *Revised and improved value of the QED tenth-order electron anomalous magnetic moment*: “ we have carried out a new numerical evaluation of the 389 integrals of Set V, which represent 6 354 Feynman vertex diagrams without lepton loops. We found that one of the integrals, called X_{024} , was given a wrong value in the previous calculation due to an incorrect assignment of integration variables. The correction of this error causes a shift of -1.26 to the Set V contribution, and hence to the tenth-order universal (mass-independent) term $7.606 (192) (\alpha/\pi)^5$ as the best estimate of the Set V contribution. ” I have entered the correction into (1.1).

References

- [1] I. K. Affleck, O. Alvarez, and N. S. Manton, “Pair production at strong coupling in weak external fields”, *Nucl. Phys. B* **197**, 509–519 (1982).
- [2] A. Ahmad, N. Ahmadiniaz, O. Corradini, S. P. Kim, and C. Schubert, “Master formulas for the dressed scalar propagator in a constant field”, *Nucl. Phys. B* **919**, 9–24 (2017).
- [3] N. Ahmadiniaz, A. Bashir, and C. Schubert, “Multiphoton amplitudes and generalized Landau-Khalatnikov-Fradkin transformation in scalar QED”, *Physical Review D* **93**, 045023s (2016).
- [4] C. Alexandrou, R. Rosenfelder, and A. W. Schreiber, “Nonperturbative mass renormalization in quenched QED from the worldline variational approach”, *Phys. Rev. D* **62**, 085009 (2000).
- [5] S. A. Anikin, O. I. Zav’yalov, and M. K. Polivanov, “Simple proof of the Bogolyubov - Parasyuk theorem”, *Theor. Math. Phys.* **17**, 1082–1088 (1973).
- [6] D. Antonov, “World-line formalism: Non-perturbative applications”, *Universe* **2**, 28 (2016).
- [7] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, “Quantum electrodynamics calculation of lepton anomalous magnetic moments: Numerical approach to the perturbation theory of QED”, *Prog. Theor. Exp. Phys.* **2012**, 01A107 (2012).
- [8] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, “Tenth-order QED contribution to the electron $g-2$ and an improved value of the fine structure constant”, *Phys. Rev. Lett.* **109**, 111807 (2012).
- [9] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, “Tenth-order electron anomalous magnetic moment: Contribution of diagrams without closed lepton loops”, *Phys. Rev. D* **91**, 033006 (2015).
- [10] T. Aoyama, M. Hayakawa, T. Kinoshita, M. Nio, and N. Watanabe, “Eighth-order vacuum-polarization function formed by two light-by-light-scattering diagrams and its contribution to the tenth-order electron $g-2$ ”, *Phys. Rev. D* **78**, 053005 (2008).
- [11] T. Aoyama, T. Kinoshita, and M. Nio, “Revised and improved value of the QED tenth-order electron anomalous magnetic moment”, *Phys. Rev. D* **97**, 036001 (2018).
- [12] N. Arkani-Hamed, T.-C. Huang, and Y.-t. Huang, *Scattering amplitudes for all masses and spins*, 2017.
- [13] S. Badger, N. E. J. Bjerrum-Bohr, and P. Vanhove, “Simplicity in the structure of QED and gravity amplitudes”, *J. High Energy Phys.* **2009**, 038 (2009).
- [14] D. Bardin, M. Bilenky, D. Lehner, A. Olchevski, and T. Riemann, “Semi-analytical approach to four-fermion production in $e+e$ -annihilation”, *Nucl. Phys. B Proc. Suppl.* **37**, 148–157 (1994).

- [15] K. Barro-Bergflödt, R. Rosenfelder, and M. Stingl, “Variational worldline approximation for the relativistic two-body bound state in a scalar model”, *Few-Body Syst.* **39**, 193–253 (2006).
- [16] J. Barrow-Green, *Poincaré and the Three Body Problem* (Amer. Math. Soc., Providence R.I., 1997).
- [17] F. Bastianelli, A. Huet, C. Schubert, R. Thakur, and A. Weber, “Integral representations combining ladders and crossed-ladders”, *J. High Energy Phys.* **2014**, 66 (2014).
- [18] R. Bermudez, L. Albino, L. X. Gutiérrez-Guerrero, M. E. Tejada-Yeomans, and A. Bashir, “Quark-gluon vertex: A perturbation theory primer and beyond”, *Phys. Rev. D* **95**, 034041 (2017).
- [19] Z. Bern and D. A. Kosower, “Efficient calculation of one-loop QCD amplitudes”, *Phys. Rev. Lett.* **66**, 1669–1672 (1991).
- [20] Z. Bern and D. A. Kosower, “The computation of loop amplitudes in gauge theories”, *Nucl. Phys. B* **379**, 451–561 (1992).
- [21] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Fields* (McGraw-Hill, 1965).
- [22] E. B. Bogomolny and Y. Kubyshin, “Asymptotic estimates for graphs with a fixed number of fermion loops in quantum electrodynamics. the choice of the form of the steepest-descent solutions”, *Sov. J. Nucl. Phys.* **34**, 853–858 (1981).
- [23] B. M. Bolotovskiy, M. A. Vasil’ev, B. L. Voronov, V. L. Ginzburg, A. V. Gurevich, N. S. Kardashev, A. A. Komar, L. V. Keldysh, A. I. Nikishov, M. A. Solov’ev, I. V. Tyutin, and A. E. Shabad, “Vladimir Ivanovich Ritus (on his eightieth birthday)”, *Phys. Usp.* **50**, 763 (2007).
- [24] E. Boos and T. Ohl, “Minimal gauge invariant classes of tree diagrams in gauge theories”, *Phys. Rev. Lett.* **83**, 480–483 (1999).
- [25] M. Borinsky, “Feynman graph generation and calculations in the Hopf algebra of Feynman graphs”, *Comput. Phys. Commun.* **185**, 3317–3330 (2014).
- [26] M. Borinsky, “Renormalized asymptotic enumeration of Feynman diagrams”, *Annals of Physics* (2017) [10.1016/j.aop.2017.07.009](https://doi.org/10.1016/j.aop.2017.07.009).
- [27] N. Brambilla, S. Eidelman, P. Foka, and et al, “QCD and strongly coupled gauge theories: challenges and perspectives”, *Eur. Phys. J. C* **74**, 2981 (2014).
- [28] D. J. Broadhurst, “Four-loop Dyson-Schwinger-Johnson anatomy”, *Phys. Lett. B* **466**, 319–325 (1999).
- [29] D. J. Broadhurst, R. Delbourgo, and D. Kreimer, “Unknotting the polarized vacuum of quenched QED”, *Phys. Lett. B* **366**, 421–428 (1996).
- [30] S. J. Brodsky and J. D. Sullivan, “*W*-boson contribution to the anomalous magnetic moment of the muon”, *Phys. Rev.* **156**, 1644–1647 (1967).
- [31] R. Carroll, “Mass-operator calculation of the electrongfactor”, *Phys. Rev. D* **12**, 2344–2355 (1975).

- [32] R. Carroll and Y.-P. Yao, “ α^3 contributions to the anomalous magnetic moment of an electron in the mass-operator formalism”, *Phys. Lett. B* **48**, 125–127 (1974).
- [33] L. Chang, Y.-X. Liu, and C. D. Roberts, “Dressed-quark anomalous magnetic moments”, *Phys. Rev. Lett.* **106**, 072001 (2011).
- [34] K. G. Chetyrkin and V. A. Smirnov, “R*-operation corrected”, *Physics Letters B* **144**, 419–424 (1984).
- [35] K. G. Chetyrkin and F. V. Tkachov, “Infrared R-operation and ultraviolet counterterms in the MS-scheme”, *Phys. Lett. B* **114**, 340–344 (1982).
- [36] I. D. Choudhury and A. Lahiri, “Anomalous chromomagnetic moment of quarks”, *Mod. Phys. Lett. A* **30**, 1550113 (2015).
- [37] I. Crump, *Graph Invariants with Connections to the Feynman Period in ϕ^4 Theory*, PhD thesis (Simon Fraser Univ., Burnaby, BC, Canada, 2017).
- [38] C. Cruz-Santiago, P. Kotko, and A. M. Stařto, “Scattering amplitudes in the light-front formalism”, *Prog. Particle Nuclear Phys.* **85**, 82–131 (2015).
- [39] P. Cvitanović, “Infra-red structure of Yang-Mills theories”, *Phys. Lett. B* **65**, 272–274 (1976).
- [40] P. Cvitanović, “Yang-Mills theory of the mass-shell”, *Phys. Rev. Lett.* **37**, 1528–1531 (1976).
- [41] P. Cvitanović, “Asymptotic estimates and gauge invariance”, *Nucl. Phys. B* **127**, 176–188 (1977).
- [42] P. Cvitanović, “Quantum chromodynamics on the mass shell”, *Nucl. Phys. B* **130**, 114–144 (1977).
- [43] P. Cvitanović, *Field theory*, tech. rep., Notes prepared by E. Gyldenkerne (Nordita, Copenhagen, 1983).
- [44] P. Cvitanović, “Chaotic Field Theory: A sketch”, *Physica A* **288**, 61–80 (2000).
- [45] P. Cvitanović, R. Artuso, R. Mainieri, G. Tanner, and G. Vattay, *Chaos: Classical and Quantum* (Niels Bohr Inst., Copenhagen, 2017).
- [46] P. Cvitanović, C. P. Dettmann, R. Mainieri, and G. Vattay, “Trace formulas for stochastic evolution operators: Weak noise perturbation theory”, *J. Stat. Phys.* **93**, 981–999 (1998).
- [47] P. Cvitanović, C. P. Dettmann, R. Mainieri, and G. Vattay, “Trace formulae for stochastic evolution operators: Smooth conjugation method”, *Nonlinearity* **12**, 939 (1999).
- [48] P. Cvitanović and T. Kinoshita, “Feynman-Dyson rules in parametric space”, *Phys. Rev. D* **10**, 3978–3991 (1974).
- [49] P. Cvitanović and T. Kinoshita, “New approach to the separation of ultraviolet and infrared divergences of Feynman-parametric integrals”, *Phys. Rev. D* **10**, 3991–4006 (1974).

- [50] P. Cvitanović and T. Kinoshita, “Sixth-order magnetic moment of the electron”, *Phys. Rev. D* **10**, 4007–4031 (1974).
- [51] P. Cvitanović, P. G. Lauwers, and P. N. Scharbach, “Gauge invariance structure of Quantum Chromodynamics”, *Nucl. Phys. B* **186**, 165–186 (1981).
- [52] P. Cvitanović, N. Søndergaard, G. Palla, G. Vattay, and C. P. Dettmann, “Spectrum of stochastic evolution operators: Local matrix representation approach”, *Phys. Rev. E* **60**, 3936–3941 (1999).
- [53] A. K. Das, J. Frenkel, and C. Schubert, “Infrared divergences, mass shell singularities and gauge dependence of the dynamical fermion mass”, *Phys. Lett. B* **720**, 414–418 (2013).
- [54] P. A. M. Dirac, “The quantum theory of the electron”, *Proc. Roy. Soc. Lond. A* **117**, 610–624 (1928).
- [55] L. J. Dixon, Calculating scattering amplitudes efficiently, in *QCD and beyond. Proceedings, Theoretical Advanced Study Institute in Elementary Particle Physics, TASI-95, Boulder, USA, June 4-30, 1995* (1996), pp. 539–584.
- [56] S. D. Drell and H. R. Pagels, “Anomalous magnetic moment of the electron, muon, and nucleon”, *Phys. Rev.* **140**, B397–B407 (1965).
- [57] G. V. Dunne and M. Krasnansky, ““Background field integration-by-parts” and the connection between one-loop and two-loop Heisenberg-Euler effective actions”, *J. High Energy Phys.* **2006**, 020 (2006).
- [58] G. V. Dunne and C. Schubert, “Two-loop self-dual Euler-Heisenberg Lagrangians (I): Real part and helicity amplitudes”, *J. High Energy Phys.* **2002**, 053 (2002).
- [59] G. V. Dunne and C. Schubert, “Two-loop self-dual Euler-Heisenberg Lagrangians (II): Imaginary part and Borel analysis”, *J. High Energy Phys.* **2002**, 042 (2002).
- [60] G. V. Dunne and C. Schubert, “Multiloop information from the QED effective Lagrangian”, *J. Phys. Conf. Ser.* **37**, 59–72 (2006).
- [61] F. J. Dyson, “Divergence of perturbation theory in Quantum Electrodynamics”, *Phys. Rev.* **85**, 631–632 (1952).
- [62] A. C. Edison and S. G. Naculich, “Symmetric-group decomposition of SU(N) group-theory constraints on four-, five-, and six-point color-ordered amplitudes at all loop orders”, *J. High Energy Phys.* **2012**, 69 (2012).
- [63] M. I. Eides, “Recent ideas on the calculation of lepton anomalous magnetic moments”, *Phys. Rev. D* **90**, 057301 (2014).
- [64] T. Epelbaum, F. Gelis, and B. Wu, “Lattice worldline representation of correlators in a background field”, *J. High Energy Phys.* **2015**, 148 (2015).
- [65] T. Epelbaum, F. Gelis, and B. Wu, “From lattice Quantum Electrodynamics to the distribution of the algebraic areas enclosed by random walks on Z^2 ”, *Ann. Inst. Henri Poincaré* **3**, 381–404 (2016).
- [66] M. Fael and M. Passera, “Positronium contribution to the electron $g-2$ ”, *Phys. Rev. D* **90**, 056004 (2014).

- [67] R. P. Feynman, “Mathematical formulation of the quantum theory of electromagnetic interaction”, *Phys. Rev.* **80**, 440–457 (1950).
- [68] R. P. Feynman, “An operator calculus having applications in Quantum Electrodynamics”, *Phys. Rev.* **84**, 108–128 (1951).
- [69] R. P. Feynman, “Slow electrons in a polar crystal”, *Phys. Rev.* **97**, 660–665 (1955).
- [70] R. P. Feynman, “The present status of Quantum Electrodynamics”, in *The Quantum Theory of Fields: Proceedings of the XII on Physics at the Univ. of Brussels* (Interscience, 1962), p. 61.
- [71] V. Fock, “Die Eigenzeit in der klassischen und in der Quantenmechanik”, *Physik. Z. Sowjetunion* **12**, Transl. Proper time in classical and quantum mechanics, 404–425 (1937).
- [72] E. S. Fradkin, “Application of functional methods in quantum field theory and quantum statistics (II)”, *Nucl. Phys. B* **76**, 588–624 (1966).
- [73] J. Frenkel, R. Meuldermans, I. Mohammad, and J. C. Taylor, “Infra-red behaviour in non-abelian gauge theories”, *Phys. Lett. B* **64**, 211–212 (1976).
- [74] H. M. Fried, *Modern Functional Quantum Field Theory: Summing Feynman Graphs* (World Scientific, Singapore, 2014).
- [75] H. M. Fried and Y. Gabellini, “Analytic, nonperturbative, almost exact QED: The two-point functions”, *Phys. Rev. D* **79**, 065035 (2009).
- [76] H. M. Fried and Y. Gabellini, “On the summation of Feynman graphs”, *Ann. Phys.* **327**, 1645–1667 (2012).
- [77] H. M. Fried and Y. Gabellini, “QED vacuum loops and vacuum energy”, *Eur. Phys. J. C* **73**, 2642 (2013).
- [78] M. P. Fry, “Nonperturbative quantization of the electroweak model’s electrodynamic sector”, *Phys. Rev. D* **91**, 085026 (2015).
- [79] P. Gaspard and D. Alonso, “ \hbar expansion for the periodic-orbit quantization of hyperbolic systems”, *Phys. Rev. A* **47**, R3468–R3471 (1993).
- [80] F. Gelis and N. Tanji, “Schwinger mechanism revisited”, *Prog. Part. Nucl. Phys.* **87**, 1–49 (2016).
- [81] E. Getzler and M. M. Kapranov, “Modular operads”, *Compositio Math.* **110**, 65–125 (1998).
- [82] H. Gies and J. Hämmerling, “Geometry of spin-field coupling on the worldline”, *Phys. Rev. D* **72**, 065018 (2005).
- [83] H. Gies and K. Langfeld, “Loops and loop clouds — a numerical approach to the worldline formalism in QED”, *Int. J. Mod. Phys. A* **17**, 966–976 (2002).
- [84] H. Gies, J. Sanchez-Guillen, and R. A. Vázquez, “Quantum effective actions from nonperturbative worldline dynamics”, *J. High Energy Phys.* **2005**, 067 (2005).

- [85] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics* (Springer, New York, 1990).
- [86] J. Henn, R. N. Lee, A. V. Smirnov, V. A. Smirnov, and A. Steinhauser, “Four-loop photon quark form factor and cusp anomalous dimension in the large- N_c limit of QCD”, *J. High Energy Phys.* **2017**, 139 (2017).
- [87] F. Herzog and B. Ruijl, “The R^* -operation for Feynman graphs with generic numerators”, *J. High Energy Phys.* **2017**, 37 (2017).
- [88] I. Huet, D. G. C. McKeon, and C. Schubert, “Euler-Heisenberg lagrangians and asymptotic analysis in 1+1 QED. Part I: Two-loop”, *J. High Energy Phys.* **2010**, 36 (2010).
- [89] I. Huet, M. R. de Trautenberg, and C. Schubert, “Asymptotic behaviour of the QED perturbation series”, in *5th Winter Workshop on Non-Perturbative Quantum Field Theory, Sophia-Antipolis*, edited by C. Schubert (2017).
- [90] I. Huet, M. R. de Trautenberg, and C. Schubert, “Multiloop Euler-Heisenberg Lagrangians, Schwinger pair creation, and the photon S-matrix”, *Russ. Phys. J.*, 1–6 (2017).
- [91] F. Jegerlehner, *The Anomalous Magnetic Moment of the Muon* (Springer, Berlin, 2017).
- [92] F. Jegerlehner and A. Nyffeler, “The muon $g - 2$ ”, *Phys. Rep.* **477**, 1–110 (2009).
- [93] S. Jia and M. R. Pennington, “How gauge covariance of the fermion and boson propagators in QED constrain the effective fermion-boson vertex”, *Phys. Rev. D* **94**, 116004 (2016).
- [94] S. Jia and M. R. Pennington, “Gauge covariance of the fermion Schwinger-Dyson equation in QED”, *Phys. Lett. B* **769**, 146–151 (2017).
- [95] S. Jia and M. R. Pennington, “Landau-Khalatnikov-Fradkin transformation for the fermion propagator in QED in arbitrary dimensions”, *Phys. Rev. D* **95**, 076007 (2017).
- [96] K. Johnson and B. Zumino, “Gauge dependence of the wave-function renormalization constant in Quantum Electrodynamics”, *Phys. Rev. Lett.* **3**, 351–352 (1959).
- [97] G. Kälin, “Cyclic Mario worlds – color-decomposition for one-loop QCD”, *J. High Energy Phys.* **2018**, 141 (2018).
- [98] N. G. van Kampen, *Stochastic Processes in Physics and Chemistry* (North-Holland, Amsterdam, 1992).
- [99] A. Katok and B. Hasselblatt, *Introduction to the modern theory of dynamical systems* (Cambridge Univ. Press, Cambridge, 1995).
- [100] T. Kinoshita, “Note on the infrared catastrophe”, *Prog. Theor. Phys.* **5**, 1045 (1950).
- [101] T. Kinoshita, “Mass singularities of Feynman amplitudes”, *J. Math. Phys.* **3**, 650–677 (1962).

- [102] T. Kinoshita, “Everyone makes mistakes—including Feynman”, *Journal of Physics G* **29**, 9 (2003).
- [103] T. Kinoshita, “Personal recollections, 1945–1960”, *Prog. Theor. Exp. Phys.* **2016**, 07B107 (2016).
- [104] T. Kinoshita and P. Cvitanović, “Sixth-order radiative corrections to the electron magnetic moment”, *Phys. Rev. Lett.* **29**, 1534–1537 (1972).
- [105] T. Kinoshita and W. B. Lindquist, “Eighth-order magnetic moment of the electron. V. Diagrams containing no vacuum-polarization loop”, *Phys. Rev. D* **42**, 636–655 (1990).
- [106] H. Kießler, “Hopf-algebraic renormalization of qed in the linear covariant gauge”, *Ann. Phys.* **372**, 159–174 (2016).
- [107] H. Kießler and D. Kreimer, “Diagrammatic cancellations and the gauge dependence of QED”, *Phys. Lett. B* **764**, 318–321 (2017).
- [108] R. Kleiss and H. Kuijf, “Multigluon cross sections and 5-jet production at hadron colliders”, *Nucl. Phys. B* **312**, 616–644 (1989).
- [109] B. Kol and R. Shir, “Color structures and permutations”, *J. High Energy Phys.* **2014**, 20 (2014).
- [110] B. Kol and R. Shir, “1-loop color structures and sunny diagrams”, *Journal of High Energy Physics* **2015**, 1–25 (2015).
- [111] C. P. Korthals Altes and E. De Rafael, “Infrared structure of non-abelian gauge theories: An instructive calculation”, *Phys. Lett. B* **62**, 320–322 (1976).
- [112] C. P. Korthals Altes and E. De Rafael, “Infrared structure of non-abelian gauge theories: Comments on perturbation theory calculations”, *Nucl. Phys. B* **125**, 275–284 (1977).
- [113] D. Kreimer, “On overlapping divergences”, *Commun. Math. Phys.* **204**, 669–689 (1999).
- [114] D. Kreimer, *Knots and Feynman diagrams* (Cambridge Univ. Press, 2000).
- [115] D. Kreimer and K. Yeats, “Recursion and growth estimates in renormalizable quantum field theory”, *Commun. Math. Phys.* **279**, 401–427 (2008).
- [116] S. Laporta, “High-precision calculation of the 4-loop contribution to the electron $g-2$ in QED”, *Phys. Lett. B* **772**, 232–238 (2017).
- [117] S. Laporta, “QED contributions to electron $g-2$ ”, *EPJ Web of Conferences* **179**, 01010 (2018).
- [118] S. Laporta and E. Remiddi, “The analytical value of the electron ($g-2$) at order α^3 in QED”, *Phys. Lett. B* **379**, 283–291 (1996).
- [119] A. Lasota and M. MacKey, *Chaos, Fractals, and Noise; Stochastic Aspects of Dynamics* (Springer, New York, 1994).
- [120] M. Laufer and P. Orland, “Metric of Yang-Mills orbit space on the lattice”, *Phys. Rev. D* **88**, 065018 (2013).

- [121] B. Lautrup, “On high order estimates in QED”, *Phys. Lett. B* **69**, 109–111 (1977).
- [122] B. E. Lautrup, A. Peterman, and E. de Rafael, “Recent developments in the comparison between theory and experiments in quantum electrodynamics”, *Phys. Rep.* **3**, 193–259 (1972).
- [123] S. L. Lebedev and V. I. Ritus, “Virial representation of the imaginary part of the Lagrange function of the electromagnetic field”, *Sov. Phys. JETP* **59**, 237 (1984).
- [124] M. J. Levine and J. Wright, “Anomalous magnetic moment of the electron”, *Phys. Rev. D* **8**, 3171–3179 (1973).
- [125] F. Maltoni, K. Paul, T. Stelzer, and S. Willenbrock, “Color-flow decomposition of QCD amplitudes”, *Phys. Rev. D* **67**, 014026 (2003).
- [126] E. Marinari, G. Parisi, and C. Rebbi, “Monte Carlo simulation of the massive Schwinger model”, *Nucl. Phys. B* **190**, 734–750 (1981).
- [127] L. C. Martin, C. Schubert, and V. M. Villanueva, “On the low-energy limit of the QED N-photon amplitudes”, *Nucl. Phys. B* **668**, 335–344 (2003).
- [128] K. Melnikov, A. Vainshtein, and M. Voloshin, “Remarks on the effect of bound states and threshold in g^{-2} ”, *Phys. Rev. D* **90**, 017301 (2014).
- [129] J. W. Mielniczuk, D. Lamm, S. Auddy, and S. R. Valluri, *The anomalous magnetic moment of a photon propagating in a magnetic field*, 2017.
- [130] A. G. Morgan, “Second order fermions in gauge theories”, *Phys. Lett. B* **351**, 249–256 (1995).
- [131] Y. Nambu, “The use of the proper time in Quantum Electrodynamics I”, *Prog. Theor. Phys.* **5**, 82–94 (1950).
- [132] T. Nieuwenhuis and J. A. Tjon, “Nonperturbative study of generalized ladder graphs in a $\phi^2\chi$ theory”, *Phys. Rev. Lett.* **77**, 814–817 (1996).
- [133] B. Penante, *On-shell methods for off-shell quantities in N=4 Super Yang-Mills: from scattering amplitudes to form factors and the dilatation operator*, 2016.
- [134] H. Poincaré, *Les Méthodes Nouvelles de la Mécanique Céleste*, For a very readable exposition of Poincaré’s work and the development of the dynamical systems theory up to 1920’s see [16]. (Guthier-Villars, Paris, 1899).
- [135] A. Prunotto, *A Homological Approach to Feynman Diagrams in the Quantum Many-Body Theory*, Corso di Laurea in Fisica, thesis, MA thesis (Università degli Studi di Torino, 2003).
- [136] A. Prunotto, W. M. Alberico, and P. Czerski, “Feynman diagrams and rooted maps”, *Open Physics* **16**, 149–167 (2013).
- [137] C. Reuschle and S. Weinzierl, “Decomposition of one-loop QCD amplitudes into primitive amplitudes based on shuffle relations”, *Phys. Rev. D* **88**, 105020 (2013).

- [138] V. I. Ritus, “A symmetry relating certain processes in 2- and 4-dimensional space-times and the value $\alpha_0 = 1/4\pi$ of the bare fine structure constant”, *J. Exp. Theor. Phys.* **102**, 582–602 (2006).
- [139] V. I. Ritus, “Duality of two-dimensional field theory and four-dimensional electrodynamics leading to a finite value of the bare charge”, *Phys. Usp.* **56**, 565 (2013).
- [140] V. I. Ritus, “On a connection of the fine structure constant with zero-point fluctuations of the electromagnetic field in vacuum”, *J. Russ. Laser Res.* **36**, 101–109 (2015).
- [141] R. Rosenfelder and A. W. Schreiber, “Polaron variational methods in the particle representation of field theory. I. General formalism”, *Phys. Rev. D* **53**, 3337–3353 (1996).
- [142] R. Rosenfelder and A. W. Schreiber, “Polaron variational methods in the particle representation of field theory. II. Numerical results for the propagator”, *Phys. Rev. D* **53**, 3354–3365 (1996).
- [143] R. Rosenfelder and A. W. Schreiber, “An Abraham-Lorentz-like equation for the electron from the worldline variational approach to QED”, *Eur. Phys. J. C* **37**, 161–171 (2004).
- [144] H. H. Rugh, “The correlation spectrum for hyperbolic analytic maps”, *Nonlinearity* **5**, 1237 (1992).
- [145] B. Ruijl, T. Ueda, J. A. M. Vermaseren, and A. Vogt, “Four-loop QCD propagators and vertices with one vanishing external momentum”, *J. High Energy Phys.* **2017**, 1–50 (2017).
- [146] M. A. Samuel, “Estimates of the eighth-order corrections to the anomalous magnetic moment of the muon”, *Phys. Rev. D* **9**, 2913–2919 (1974).
- [147] M. Schäfer, I. Huet, and H. Gies, “Worldline numerics for energy-momentum tensors in Casimir geometries”, *Journal of Physics A: Mathematical and Theoretical* **49**, 135402 (2016).
- [148] M. G. Schmidt and C. Schubert, “Multiloop calculations in the string-inspired formalism: The single spinor loop in QED”, *Phys. Rev. D* **53**, 2150–2159 (1996).
- [149] M. G. Schmidt and I. O. Stamatescu, “Matter determinants in background fields using random walk world line loops on the lattice”, *Mod. Phys. Lett. A* **18**, 1499–1515 (2003).
- [150] M. G. Schmidt and I.-O. Stamatescu, “Determinant calculations using random walk worldline loops”, *Nucl. Phys. B Proc. Suppl.* **119**, Proceedings of the XXth International Symposium on Lattice Field Theory, 1030–1032 (2003).
- [151] O. Schnetz, *Numbers and functions in Quantum Field Theory*, 2016.
- [152] C. Schubert, “Perturbative quantum field theory in the string-inspired formalism”, *Phys. Rep.* **355**, 73–234 (2001).

- [153] C. Schubert, “Lectures on the worldline formalism”, in *School of Spinning Particles in Quantum Field Theory: Worldline Formalism, Higher Spins and Conformal Geometry*, edited by C. Schubert (2012).
- [154] T. Schuster, “Color ordering in QCD”, *Phys. Rev. D* **89**, 105022 (2014).
- [155] M. D. Schwartz, *Quantum Field Theory and the Standard Model* (Cambridge Univ. Press, Cambridge, 2013).
- [156] J. Schwinger, “On Quantum-Electrodynamics and the magnetic moment of the electron”, *Phys. Rev.* **73**, 416–417 (1948).
- [157] J. Schwinger, “On gauge invariance and vacuum polarization”, *Phys. Rev.* **82**, 664–679 (1951).
- [158] J. Schwinger, “On the Green’s functions of quantized fields. I”, *Proc. Natl. Acad. Sci. USA* **37**, 452–455 (1951).
- [159] J. Schwinger, “On the Green’s functions of quantized fields. II”, *Proc. Natl. Acad. Sci. USA* **37**, 455–459 (1951).
- [160] E. Seiler and I. O. Stamatescu, “A note on the loop formula for the fermionic determinant”, *J. Phys. A* **49**, 335401 (2016).
- [161] Y. Simonov, “Relativistic path integral and relativistic Hamiltonians in QCD and QED”, *Phys. Rev. D* **88**, 025028 (2013).
- [162] Y. A. Simonov, “QED spectra in the path integral formalism”, *Phys. Rev. D* **90**, 013013 (2014).
- [163] Y. A. Simonov and J. Tjon, “The Feynman–Schwinger representation in QCD”, *Ann. Phys.* **300**, 54–87 (2002).
- [164] A. A. Slavnov, “Ward identities in gauge theories”, *Theor. Math. Phys.* **10**, 99–104 (1972).
- [165] V. A. Smirnov and K. G. Chetyrkin, “R* operation in the minimal subtraction scheme”, *Theoretical and Mathematical Physics* **63**, 462–469 (1985).
- [166] C. M. Sommerfield, “The magnetic moment of the electron”, *Ann. Phys.* **5**, 26–57 (1958).
- [167] N. Søndergaard, G. Palla, G. Vattay, and A. Voros, “Asymptotics of high order noise corrections”, *J. Stat. Phys.* **101**, 385–395 (2000).
- [168] E. R. Speer, “Analytic renormalization”, *J. Math. Phys.* **9**, 1404–1410 (1968).
- [169] M. J. Strassler, “Field theory without Feynman diagrams: One-loop effective actions”, *Nucl. Phys. B* **385**, 145–184 (1992).
- [170] I. M. Suslov, “High orders of perturbation theory. Are renormalons significant?”, *Sov. Phys. JETP* **89**, 197–207 (1999).
- [171] G. ’t Hooft, “A planar diagram theory for strong interactions”, *Nucl. Phys. B* **72**, 461–473 (1974).
- [172] Y. Takahashi, “On the generalized Ward identity”, *Nuovo Cim.* **6**, 371 (1957).
- [173] J. C. Taylor, “Ward identities and charge renormalization of the Yang-Mills field”, *Nucl. Phys. B* **33**, 436–444 (1971).

- [174] I. Todorov, *From Euler's play with infinite series to the anomalous magnetic moment*, 2018.
- [175] G. Torgrimsson, C. Schneider, J. Oertel, and R. Schützhold, "Dynamically assisted Sauter-Schwinger effect — non-perturbative versus perturbative aspects", *J. High Energy Phys.* **2017**, 1–26 (2017).
- [176] S. R. Valluri, U. D. Jentschura, and D. R. Lamm, The study of the Heisenberg-Euler Lagrangian and some of its applications, in *Aip conf. proceed.* Vol. 687 (2003), pp. 203–222.
- [177] R. Vilela Mendes, "A consistent measure for lattice yang–mills", *Inter. J. Mod. Phys. A* **32**, 1750016 (2017).
- [178] S. A. Volkov, "Subtractive procedure for calculating the anomalous electron magnetic moment in QED and its application for numerical calculation at the three-loop level", *J. Exp. Theor. Phys.* **122**, 1008–1031 (2016).
- [179] S. A. Volkov, "New method of computing the contributions of graphs without lepton loops to the electron anomalous magnetic moment in QED", *Phys. Rev. D* **96**, 096018 (2017).
- [180] J. C. Ward, "An identity in Quantum Electrodynamics", *Phys. Rev.* **78**. 182 (1950).
- [181] K. Yeats, *A Combinatorial Perspective on Quantum Field Theory* (Springer, New York, 2017).
- [182] D. R. Yennie, S. C. Frautschi, and H. Suura, "The infrared divergence phenomena and high-energy processes", *Ann. Phys.* **13**, 379–452 (1961).
- [183] D. Zeppenfeld, "Diagonalization of color factors", *Inter. J. Mod. Phys. A* **3**, 2175–2179 (1988).

Chapter 2

Learning worldline QFT

2.1 Quantum Field Theory Spring 2018 Report

Guopeng Xu <guopengxu@gatech.edu>
term paper for Spring / Summer 2018 QFT self-study course.

The goal. Take a narrow path, learn enough Quantum Field Theory (QFT) (and Green's functions and such, but no more) to be able to digest sect. 1.3 *Worldline formalism* and check James sect. 1.3.2 *Electron magnetic moment in worldline formalism* calculations.

What is a gauge theory? According to [6] there are two types of theories that can be called 'gauge theories', the Yang-Mills theories and constrained Hamiltonian theories.

Summary In order to get correct predictions from non-Abelian field theories, which are susceptible to large number of gauge copies, we need to choose a representative of each gauge orbit.

2.2 Morelia Worldline Formalism course

2017-07-03 Predrag First Morelia discussion (all errors are mine):

Christian is inclined to compute (g-2) starting with their two-field spinor QED Bern-Kosower formula. In that formulation all photons are born equal; one of them is kept as the external field (the seagull vertex, or the k^μ coefficient) is the magnetic moment $\sigma_{\mu\nu}$), and the rest are contracted pairwise in all possible ways. In the quenched case, this yields one gauge invariant set, the self-energy set of sect. 1.2.3.

James would like to start with their electron propagator in constant external field, and keep the term linear in the external field. I vastly prefer that, because it

should be possible to distinguish in- and out-legs, and the three kinds of N -photon propagators that yield the minimal gauge sets.

I probably need to get through the proof of gauge invariance with them.

2017-07-04 Predrag Morelia Schubertiad day 1: the lecture written up in Schubert [15] 2012 *Lectures on the worldline formalism*, sects. 1.4 *Gaussian integrals* and 1.5 *The N -photon amplitude*.

Christian was right. One has to start with scalar QED one-loop effective action to understand the Bern-Kosower type master formulas. That yields a loop with any number of photons attached, each photon vertex carrying a 1D proper time Green's function. This could be computed by usual math methods techniques for computing Green's functions, but they find it useful for reasons that will be understood later to compute it as a sum of Fourier modes. The marginal modes (4 space-time translations) are fixed in the Gauss way, by shifting the origin to loops center of mass (i.e., different symmetry reduction for each loop).

We then separate the integration over x_0 , thus reducing the path integral to an integral over the relative coordinate q :

$$x^\mu(\tau) = x_0^\mu(\tau) + q^\mu(\tau), \quad (2.1)$$

with the relative coordinate q periodic and satisfying constraint

$$\int_0^T d\tau q^\mu(\tau) = 0. \quad (2.2)$$

In the symmetry-reduce q -space the zero-mode integral then yields the energy-momentum conservation δ function. The 1D Laplacian $M = -d^2/d\tau^2$ has positive eigenvalues (the usual k^2 Fourier modes), (do the exercise!)

$$\det M = (4T)^D, \quad (2.3)$$

and the bosonic Green's function of $-\frac{1}{2} \frac{d^2}{d\tau^2}$ in the symmetry-reduced space is

$$G_B^c(\tau, \tau') = 2 \langle \tau | \left(\frac{d^2}{d\tau^2} \right)^{-1} | \tau' \rangle = |\tau - \tau'| - \frac{(\tau, \tau')^2}{T} - \frac{T}{6}. \quad (2.4)$$

The first derivative \dot{G} has a sign function, and \ddot{G} has a $\delta(\tau - \tau')$ (because of the translation invariance, $d/d\tau$ can always be taken to act on the left variable τ).

This results in a Bern-Kosower [7] type master formula

$$\begin{aligned} \Gamma_{\text{scal}}[k_1, \varepsilon_1; \dots; k_N, \varepsilon_N] = & \quad (2.5) \\ & (-ie)^N (2\pi)^D \delta(\sum k_i) \int_0^\infty \frac{dT}{T} (4\pi T)^{-\frac{D}{2}} e^{-m^2 T} \prod_{i=1}^N \int_0^T d\tau_i \\ & \times \exp \left\{ \sum_{i,j=1}^N \left[\frac{1}{2} G_{Bij} k_i \cdot k_j - i \dot{G}_{Bij} \varepsilon_i \cdot k_j + \frac{1}{2} \ddot{G}_{Bij} \varepsilon_i \cdot \varepsilon_j \right] \right\} \Big|_{\text{lin}(\varepsilon_i)} \end{aligned}$$

for the one-loop N -photon amplitude in scalar QED, with photon momenta k_i and polarization vectors ε_i . m denotes the mass, e the charge and T the total proper time of the scalar loop particle.

This method is not applicable to open fermion lines.

A scary realisation. This is a special case of our general, nonlinear, arbitrary order interaction vertex smooth conjugacy calculation of sect. 1.6. In other words, our calculation is not just for scalar theory in vacuum, it is for any nonconstant background, and any order of interaction, i.e., inter alia general relativity. As we start from a saddlepoint (classical periodic solution) that is not translation invariant, we do not have to worry about fixing the marginal modes - there are none.

I fear having to explain the smooth conjugacy method to civilians.

2017-07-05 Predrag Morelia Schubertiad day 2: the $N = 2$ photon legs case is worked out in Schubert [15] 2012 lectures sect. 1.6 *The vacuum polarization*.

Even though the result is strictly zero (by Furry theorem, or by time reversal of odd number of G functions), $N = 3$ is useful to start understanding how integrations by parts work.

For N photon legs, see sect. 2.5 *Integration-by-parts and the replacement rule* and Ahmadiiaz, Schubert and Villanueva [4] *String-inspired representations of photon/gluon amplitudes*, arXiv:1211.1821: “The Bern-Kosower rules provide an efficient way for obtaining parameter integral representations of the one-loop N -photon/gluon amplitudes involving a scalar, spinor or gluon loop, starting from a master formula and using a certain integration-by-parts (“IBP”) procedure. Strassler observed that this algorithm also relates to gauge invariance, since it leads to the absorption of polarization vectors into field strength tensors. Here we present a systematic IBP algorithm that works for arbitrary N and leads to an integrand that is not only suitable for the application of the Bern-Kosower rules but also optimized with respect to gauge invariance. In the photon case this means manifest transversality at the integrand level, in the gluon case that a form factor decomposition of the amplitude into transversal and longitudinal parts is generated naturally by the IBP, without the necessity to consider the nonabelian Ward identities. Our algorithm is valid off-shell, and provides an extremely efficient way of calculating the one-loop one-particle-irreducible off-shell Green’s functions (“vertices”) in QCD. In the abelian case, we study the systematics of the IBP also for the practically important case of the one-loop N -photon amplitudes in a constant field.”

2017-07-06 Predrag Morelia Schubertiad day 3:

Idrish Huet explained to me how the numerical Monte-Carlos of worldline path integrals work.

2017-07-05 Christian says there is a relevant new arXiv from some Vietnamese authors, but I couldn’t find it.

2017-07-07 Predrag Morelia Schubertiad day 4:

2017-07-09 5:22 am Predrag had a panic attack that we'll never get started on (g-2). Forgot all about going to Patzquaro, spent entire Sunday writing up the new sect. 1.1 *Electron magnetic moment* and sect. 1.3.2 *Electron magnetic moment in worldline formalism*.

2017-07-10 Predrag Morelia Schubertiad day 5: handwritten notes only.

2017-07-11 Predrag Morelia Schubertiad day 6:

Christian assigned homework: reformulate the magnetic moment vertex operator (1.3), projections (1.4) and (1.5) in configuration coordinates.

2017-07-12 Predrag Morelia Edwardsiad day 7: **Scalar QED open lines**. Following Ahmadinia, Bashir and Schubert [1] 2016 *Multiphoton amplitudes and generalized Landau-Khalatnikov-Fradkin transformation in scalar QED*, [arXiv:1511.05087](#), and Ahmadinia, Bastianelli and Corradini [2] *Dressed scalar propagator in a non-Abelian background from the worldline formalism*, [arXiv:1508.05144](#).

Worldline is sum of $N!$ photon insertions, spinor indices and gauge transformations at the endpoints. In 1950 Feynman [11] gave the scalar QED worldline integral (1.17). Note dT for the open line, as opposed to dT/T for the closed loop: due to einbein gauge fixing that leads to different Fadeev-Popov for the loop than for the line. The worldline Green's function is now (compare to (2.4))

$$\begin{aligned} \Delta(\tau_1, \tau_2) &= \frac{1}{2}|\tau_1 - \tau_2| - \frac{1}{2}(\tau_1 - \tau_2) + \frac{\tau_1\tau_2}{T} \\ &= \frac{1}{2}(G_B(\tau_1, \tau_2) - G_B(\tau_1, 0) - G_B(0, \tau_2) + G_B(0, 0)) . \end{aligned} \quad (2.6)$$

2017-07-13 Predrag Morelia Edwardsiad day 8: handwritten notes only.

2017-07-14 Predrag Morelia Edwardsiad day 9:

Wrote down the master formula for $S_{(N)}^{xx'}$ and its kernel $K_{(N)}^{xx'}$. Rewrote it in momentum space. Verified that $N = 0$ generates the free propagator. Checked the $N = 1$ vertex. The hardest was computing the fermion self-energy, in terms of 2 dimensional regularization hypergeometric functions. Currently can compare to Davydychev [10] only numerically.

Read also Davydychev [9] *Geometrical methods in loop calculations and the three-point function*

2018-06-10 Predrag Les Houches Edwardsiad day 10:

James has an open-line Dirichlet Green's function that should suffice to compute (g-2). Predrag is sceptical, thinks that Dirichlet Green's function is a bad choice, as it breaks the translational invariance. Would prefer some periodic formulation, where a periodic space box is taken off to infinity, than put on the mass-shell either by some absorptive cut (yielding (1.5)) or amputated and renormalized by $\sqrt{Z_2}$ as in LSZ formulation.

I thought that in the quenched QED worldline formulation, the one-particle reducible graphs renormalize (g-2) contributions (my talk `reducesymm/presentations/LesHouch18/finiteQED.tex`) but my N -photon formulation of the QED vertex overcounts counter-terms, or, as Magnea says in his handwritten [lecture 2](#), “one-particle reducible graphs should be included only once.”

2.3 Xu wordline formalism notes

2018-01-25 Predrag to Guopeng Created sect. [2.3](#) for you to write your QFT study notes in (to James: they are both undergraduate students, and they are expected to form a study group. Guopeng is from Jilin University, has gone through the first part of Peskin and Schroeder [[14](#)]).

2018-01-08 Christian Schubert that’s great the Guopeng wants to learn the worldline formalism. The problem is that on the part that interests you most, the open fermion line, we still have not written up anything intelligible. But James is in the process of writing up his part of the lectures that we gave for you, which is going to be incorporated into the lecture notes that I have on the web with Olindo Corradini. So maybe for the time being Guopeng might want to work through the lectures as they are, and hopefully by the time he is through with this, James’ fermion line part may already be in shape. Any questions, we are available.

2018-01-09 James Edwards It’s great to hear that Guopeng is getting involved in the worldline formalism. Christian’s suggests that you begin with the Olindo Corradini and Christian Schubert and notes [[8](#)], [arXiv:1512.08694](#). 2012 notes by Schubert [[15](#)], ([click here](#)) may be also helpful (Predrag: I think the 2012 notes are included in the entirety into the Corradini and Schubert notes [[8](#)]). However, you might find parts of Corradini 2012 lectures useful, ([click here](#))).

In the meantime, I will write up my notes on open fermion lines and share them with you as soon as possible. Needless to say, if Guopeng has any questions he is most welcome to get in touch with us to discuss things in more detail.

2018-01-16 James Would Guopeng like to get involved in some ongoing calculations pertaining to g-2? Our worldline representation of the fermion propagator provides a new technique for attacking this problem and there has already been some progress in calculating some of the ingredients for the 3 loop contribution, but we would also like to look at 1+ loop rainbow diagrams.

To achieve this, we would study the open fermion line with a low energy photon attached along with a certain number of virtual photon loops. Now we know how to deal with the fermion line we have a good idea of how to calculate these quantities in a particularly efficient way (that automatically picks out the term linear in k with the right gamma matrix structure to allow for extraction of the structure constant we seek). We expect to be able to streamline the calculation and

maybe even earn some advantages in the limit of large numbers of virtual photons. In principle, this can be extended to the case of a constant electromagnetic background too.

We can begin working by checking the one loop contribution as a warm-up (I have notes on this already) before looking at the 2-loop rainbow. This requires familiarity with the scalar propagator in the worldline approach before looking at open spinor lines and the coupling to external photons. The latter remain as my job to write up our notes from Predrag's visit, and I will strive to do this asap.

2018-02-13 Predrag I find discrete lattice problems very useful in understanding concepts in QFT. Study ChaosBook.org/FieldTheory/postscript.html Chapter 1 Lattice field theory. You will learn that Laplacian generates all walks on a lattice (making it easier to understand path integrals), that a Green's function is the propagator (sum over all walks), that the mysterious $e^{iET/\hbar}$ and $e^{ip\cdot x/\hbar}$'s of Quantum Mechanics are roots of unity (a consequence of time and space translation invariance), and how that enables you to compute the propagators / Green's functions.

All of the above can be then continued in the small lattice spacing limit into things you accept on faith when you read QFT textbook. But here it is just matrices and vectors, and you can check every step.

Do not use much time on the discrete lattice formalism, as we shall not use it in the current project.

2018-02-18 Predrag To see whether you have actually understood Green's function on the level needed for our term's goals, see whether you can follow worldline Green's function calculations following eq. (1.22), eq. (2.6) above, eqs. (2.35) and (2.41) in Corradini and Schubert notes [8].

2018-03-27 Guopeng to James I think I have understood the material "Spinning particles in QM and QFT" [8]. I skipped the 1.4 section and 2.5. If you could give me the calculation, I can go through it.

2018-03-28 James Great to hear that you have gone through the material from the notes. At this stage, I don't think you need to worry about the gravitational case in section 1.4 and it's true that initially we are more concerned with Abelian QED (that said, I would recommend that you do take a look at section 2.5.1 when you have a chance, just because it gives the simplest extension of the worldline formalism to the non-Abelian case). These days we have some more powerful techniques to deal with colour degrees of freedom, but it's good to see the simpler presentation in that section).

I understand that you were referring to the material on open lines in the worldline approach, which is of course relevant to the g-2 calculation. I have almost written up my notes from Predrag's visit and hope to send them to you by the end of the week; I hope you don't mind being an "alpha tester" for these notes as one of the first people to go through them! I would encourage you to go through them

critically in case any typos have crept in. At the very least, I will send you the vacuum propagators for the scalar and spinor particle this week.

In the meantime, do let us know if you have any questions about the notes or exercises so far. If you're eager to get started, I would suggest that you review the path integral calculation of quantum mechanical kernels (i.e. matrix elements of the form $K(x, y; T) = \langle y | \exp[-iTH] | x \rangle$) including electromagnetic interactions that will be used repeatedly in the open line calculations. For example, sects. 2.18 to 2.20 of Kleinert [12] path integral textbook has full details.¹

2018-03-31 Predrag to Guopeng - time to start LaTeXing Skip section 2.5.1 (extension of the worldline formalism to the non-Abelian case) in Corradini and Schubert notes [8], we are very short on time. Focus on QED. I believe you have gone through the path integral calculation of quantum mechanical kernel.

Instead, please write up your derivation of the Green's functions (1.35) and (2.15), then (1.41), here, in this text, in LaTeX, with all signs and prefactors correct. If you have time before James' notes arrive, going through sect. 2.4 *The vacuum polarization* might be good - that connects these mysterious worldline Green's functions to the standard Feynman integrals like those you see in Peskin and Schroeder.

2018-04-16 Predrag to Guopeng Go through Stone and Goldbart [16], *Mathematics for Physics: A Guided Tour for Graduate Students*, Chapter 5 *Green Functions*. A pre-publication draft can be found [here](#).

It is all about 1-dimensional Green's functions - understanding material related to kind of function you need for worldline formalism should be helpful to you.

Stone and Goldbart [16] is an advanced summary where you will find almost everything one needs to know. More pedestrian and perhaps easier to read is Arfken and Weber [5].

I like Mathews and Walker [13], based on lectures by Richard Feynman at Cornell University. You can download it from [here](#). However, I'm not sure it will help you understand the problem at hand.

The above examples are the usual way students are taught Green's functions. I personally find deriving them as continuum limits of lattice formulations ([click here](#)) the most insightful and easiest to understand. My derivation is for periodic boundary conditions. You will have to also understand other kinds of boundary conditions that arise in your project.

2018-07-12 Guopeng Added the Edwards reference Ahmadiniaz *et al.* [3] *One-particle reducible contribution to the one-loop spinor propagator in a constant field*

2018-07-12 Guopeng, Predrag Added the Edwards / Cvitanović June 2018 whiteboard notes, [reducesymm/guopeng/](#):

¹Predrag: 2018-04-16 I have a copy of Kleinert [12]

Mrenorm.pdf The perturbative expansions for the magnetic moment anomaly (1.6).

M4.pdf

GammaVertex.pdf

WardIds.pdf

worldLprop.pdf

M4.pdf

aSums.pdf The next five files are gauge sets $a_{km'm}$ defined in (1.10):

a111.pdf

a111Gset.pdf

a111old.pdf

a120.pdf

a210.pdf

lloopPhi3.pdf is an outline of Edwards purported 1-loop calculation.

2018-07-13 Guopeng working through Edwards' *lloopPhi3.pdf* calculation sketch:

The note starts out with the N photon insertions propagator for a charged scalar field in external field (1.21) ²

$$\begin{aligned} N \text{ photon ins. fig} &= \int_0^\infty dT e^{-m^2 T} \\ &= (ie)^N \int_0^\infty dT e^{-m^2 T} \int_0^T d\tau_1 \cdots \int_0^T d\tau_N \\ &\quad \times \int_{x(0)=y}^{x(T)=x} \mathcal{D}x e^{i \sum_{i=1}^N k_i \cdot x(\tau_i)} e^{-\int_0^T d\tau \frac{1}{4} \dot{x}^2}. \end{aligned} \quad (2.7)$$

The expression $\# / [(x_a - x_b)^2]^{D/2-1}$ can be obtained by change of variable $u = 1/T$, and evaluating the Gaussian integral

$$\int_0^\infty \frac{dT}{(4\pi T)^{D/2}} e^{-(x_a - x_b)^2 / 4T}.$$

The Ingredient #1:

^{3 4} I calculate this by factorizing the Laplacian,

$$\begin{aligned} \text{Det} \left(-\frac{1}{4} \frac{\partial^2}{\partial \tau^2} + \frac{B_{ab}}{4T} \right) &= \text{Det} \left[-\frac{1}{4} \frac{\partial^2}{\partial \tau^2} \left(1 - \left(\frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{T} \right) \right] \\ &= \text{Det} \left(-\frac{1}{4} \frac{\partial^2}{\partial \tau^2} \right) \text{Det} \left(1 - \left(\frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{T} \right). \end{aligned} \quad (2.8)$$

²Guopeng 2018-07-29: [A confusing point about the master formula at the top of the *lloopPhi3.pdf* calculation: it seems like there is no linear part of the polarization vector. Where have those vectors gone?](#)

³Guopeng 2018-07-29: I wonder if there should be a T in the denominator of the derivative term, because I can't find the extra T in the eq. (2.11) of "Spinning particles in the QM and QFT" [8].

⁴James 2018-07-29: It is probable that in the notes you refer to the authors had made a rescaling of the interval $[0, T]$ to the unit interval, which induces a $1/T$ for the derivative term.

where $\text{Det} \left(-\frac{1}{4} \frac{\partial^2}{\partial \tau^2} \right) = (4\pi T)^{-\frac{D}{2}}$.⁵ As for the $\text{Det} \left(1 - \left(\frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{T} \right)$, it would appear that^{6 7}

$$-\left(\frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{T} = \frac{G_{Bab}}{T},$$

so that the determinant could be $(1 + \frac{G_{Bab}}{T})^{-\frac{D}{2}}$, where G_{Bab} is the loop Green's function. As

$$B_{ab}(\tau_1, \tau_2) = [\delta(\tau_a - \tau_1) - \delta(\tau_b - \tau_1)][\delta(\tau_a - \tau_2) - \delta(\tau_b - \tau_2)],$$

$-\left(\frac{\partial^2}{\partial \tau_1^2} \right)^{-1} \frac{B_{ab}}{T}$ is proportional to $\tau - \frac{1}{2}(\tau_a + \tau_b)$, since

$$\begin{aligned} \frac{\partial^2}{\partial \tau_1^2} |\tau_1 - \tau_2| &= 2\delta(\tau_1 - \tau_2) \\ |\tau_a - \tau| - |\tau_b - \tau| &= 2\tau - (\tau_a + \tau_b). \end{aligned} \quad (2.9)$$

8

The Ingredient #2:

$$\Delta_{(1)}(\tau_1, \tau_2 | \tau_a, \tau_b) := \langle \tau_1 | \frac{1}{-\frac{1}{4T} \frac{\partial^2}{\partial \tau^2} + \frac{B_{ab}}{4T}} | \tau_2 \rangle. \quad (2.10)$$

For $\langle \tau_1 | \frac{B_{ab}}{4T} | \tau_2 \rangle$, we have

$$\begin{aligned} &\langle \tau_1 | B_{ab}(\tau_1, \tau_2) | \tau_2 \rangle \\ &= \langle \tau_1 | [\delta(\tau_a - \tau_1) - \delta(\tau_b - \tau_1)][\delta(\tau_a - \tau_2) - \delta(\tau_b - \tau_2)] | \tau_2 \rangle \\ &= \int_0^\infty d\tau \langle \tau_1 | [\delta(\tau_a - \tau_1) - \delta(\tau_b - \tau_1)][\delta(\tau_a - \tau_2) - \delta(\tau_b - \tau_2)] | \tau \rangle \langle \tau | \tau_2 \rangle \\ &= \delta(\tau_1 - \tau_2) ([\delta(\tau_a - \tau_1) - \delta(\tau_b - \tau_1)][\delta(\tau_a - \tau_2) - \delta(\tau_b - \tau_2)]) \end{aligned} \quad (2.11)$$

since $\langle \tau_1 | \tau \rangle = \delta(\tau_1 - \tau)$, $\langle \tau | \tau_2 \rangle = \delta(\tau - \tau_2)$ and

$$\int_0^\infty d\tau \delta(\tau_1 - \tau) \delta(\tau - \tau_2) = \delta(\tau_1 - \tau_2),$$

but this calculation doesn't match the result in the *loopPhi3.pdf* note.

However, note that when we calculate $\text{Det} \left(1 - \left(\frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{T} \right)$, we assume that $-\left(\frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{T} = \frac{G_{Bab}}{T}$,⁹ so the matrix is diagonal matrix, that's why we have

$$\text{Det} \left(1 - \left(\frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{T} \right) = \left(1 + \frac{G_{Bab}}{T} \right)^{-\frac{D}{2}}$$

⁵Predrag: 2018-08-04 to Guopeng: Have you gone through exercise of evaluating (2.3)? If so, write it up.

⁶James 2018-07-29: This is not correct! See Chris' review eq. (8.9) (you are missing some delta functions).

⁷Guopeng 2018-08-03: What's the eq. (8.9) in Chris's review that you refer to? I have looked at tree level notes and "Spinning particles in QM and QFT" [8], and I find no eq. (8.9).

⁸Guopeng 2018-07-29: I don't know why this is equal to the loop Green's function.

⁹Guopeng 2018-07-29: I don't know how to prove this.

but in terms of the result in James calculation,

$$\langle \tau_1 | B_{ab}(\tau_1, \tau_2) | \tau_2 \rangle = \frac{(\Delta_{1a} - \Delta_{1b})(\Delta_{a2} - \Delta_{b2})}{\bar{T} + G_{Bab}}$$

when $\tau_1 \neq \tau_2$, the matrix element should be zero if it is diagonal, so the matrix shouldn't be diagonal, then how is the determinant to be calculated?

2018-07-30 Predrag Best to make intermediate calculation questions into footnotes, which you then comment out as you correct (or expand upon) the steps in the calculation as you fix them, so the resulting blog is the clean, correct calculation.

Also, one always has a space after ‘,’ and ‘.’ but not a space before them.

2018-08-01 Predrag to Guopeng: A crucial ingredient in any research is keeping up with literature. I did not write down the lecture notes for **2017-07-12 Predrag** Morelia Edwardsiad day 7 (above), but presumably the reference Ahmadiniaz, Bashir and Schubert [1] preceding (2.6) teaches you how to do one-loop vertex corrections for scalar QED. Though I have to admit that I do not see Edwards' Green's functions with different boundary conditions there. So just give a quick read to the two articles, and than maybe we go in depth in some calculation in them.

Possibly references listed under **2017-07-14 Predrag** are also relevant.

2018-08-03 Guopeng In *lloopPhi3.pdf* note the $\Delta_{(1)}$ is defined as in (2.10). If I extract $-\frac{1}{4T} \frac{\partial^2}{\partial \tau^2}$, and expand the latter term,¹⁰ I obtain

$$\begin{aligned} \Delta_{(1)}(\tau_1, \tau_2 | \tau_a, \tau_b) &= \langle \tau_1 | \left[-\frac{1}{4T} \frac{\partial^2}{\partial \tau^2} \right]^{-1} \left[1 + \left(-\frac{1}{4T} \frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{4\bar{T}} \right]^{-1} | \tau_2 \rangle \\ &\simeq \langle \tau_1 | \left[-\frac{1}{4T} \frac{\partial^2}{\partial \tau^2} \right]^{-1} \left[1 - \left(-\frac{1}{4T} \frac{\partial^2}{\partial \tau^2} \right)^{-1} \frac{B_{ab}}{4\bar{T}} \right] | \tau_2 \rangle \\ &= \langle \tau_1 | \left[\left(-\frac{1}{4T} \frac{\partial^2}{\partial \tau^2} \right)^{-1} - \frac{B_{ab}}{4\bar{T}} \right] | \tau_2 \rangle. \end{aligned} \quad (2.12)$$

Using

$$\langle \tau_1 | \left(-\frac{1}{4T} \frac{\partial^2}{\partial \tau^2} \right)^{-1} | \tau_2 \rangle = \Delta(\tau_1, \tau_2)$$

I get

$$\langle \tau_1 | -\frac{B_{ab}}{4\bar{T}} | \tau_2 \rangle = \frac{(\Delta_{1a} - \Delta_{1b})(\Delta_{a2} - \Delta_{b2})}{\bar{T} + G_{Bab}},$$

so it appears again that we need to know how to prove the latter equation. The reason why I expand the latter term is that I assume the deviation from the classical path is small. But I could be wrong.

Currently I am looking through Ahmadiniaz, Bashir and Schubert [1] and preparing my TOEFL exam:)

¹⁰Predrag: 2018-08-04 to Guopeng: I hope we are not allowed any such “small deviation” approximations, and that Edwards' Green's functions are exact. But I have not gone through *lloopPhi3.pdf* myself.

2018-08-14 Guopeng A small handwriting mistake is founded. In treelevel note, there misses a charge in the right side of equation 1.97 in the second term.

Some thoughts or questions about worldline formalism. When we derive the master formula for scalar QED, we assume that the electromagnetic field is composed of N photons, so we can use this formula to describe a scalar particle (like a proton or electron) interact with N photons. So when $N = 0$, it describe a free scalar particle. When $N = 1$, it describe a scalar particle emit one photon or absorb one photon, in this case, I wonder whether it satisfy the Special Relativity. For an electron emit a photon, if we observe this process in the center of mass frame with respect to ingoing electron, then it will violate the energy conservation law. For $N = 2$, a scalar particle emit(absorb) a photon and then absorb(emit) a photon at somewhere else, if the two photon is two different photons (I mean the source is different), then this describe Compton scattering process. If the two photon is the same one (for instance, emitted first then absorbed by the same electron), this describe self-energy process, in this case, we take the replacement

$$\epsilon_\mu \epsilon_\nu \longrightarrow \int \frac{d^D k}{(2\pi)^D} \frac{\eta_{\mu\nu}}{k^2}$$

in master formula. This is obtained from "sew" two photon together.

In 1loopPhi3 calculation. The extra term behind the master formula

$$\int_0^T d\tau_a \int_0^T d\tau_b \frac{\#}{[(x(\tau_a) - x(\tau_b))^2]^{D/2-1}}$$

This act like the photon vector operator, thus we add this term.

References

- [1] N. Ahmadinia, A. Bashir, and C. Schubert, "Multiphoton amplitudes and generalized Landau-Khalatnikov-Fradkin transformation in scalar QED", *Physical Review D* **93**, 045023s (2016).
- [2] N. Ahmadinia, F. Bastianelli, and O. Corradini, "Dressed scalar propagator in a non-abelian background from the worldline formalism", *Phys. Rev. D* **93**, 025035 (2016).
- [3] N. Ahmadinia, F. Bastianelli, O. Corradini, J. P. Edwards, and C. Schubert, "One-particle reducible contribution to the one-loop spinor propagator in a constant field", *Nucl. Phys. B* **924**, 377–386 (2017).
- [4] N. Ahmadinia, C. Schubert, and V. M. Villanueva, "String-inspired representations of photon/gluon amplitudes", *J. High Energy Phys.* **2013**, 132 (2013).
- [5] G. B. Arfken and H. J. Weber, *Mathematical Methods for Physicists: A Comprehensive Guide*, 6th ed. (Academic, New York, 2005).
- [6] G. Belot, "Symmetry and gauge freedom", *Stud. Hist. Philos. M. P.* **34**, 189–225 (2003).

- [7] Z. Bern and D. A. Kosower, “Efficient calculation of one-loop QCD amplitudes”, *Phys. Rev. Lett.* **66**, 1669–1672 (1991).
- [8] O. Corradini and C. Schubert, *Spinning particles in quantum mechanics and quantum field theory*, 2015.
- [9] A. I. Davydychev, “Geometrical methods in loop calculations and the three-point function”, *Nucl. Instrum. Methods Phys. Res., Sect. A* **559**, 293–297 (2006).
- [10] A. I. Davydychev, P. Osland, and L. Saks, “Quark-gluon vertex in arbitrary gauge and dimension”, *Phys. Rev. D* **63**, 014022 (2000).
- [11] R. P. Feynman, “Mathematical formulation of the quantum theory of electromagnetic interaction”, *Phys. Rev.* **80**, 440–457 (1950).
- [12] H. Kleinert, *Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets* (World Scientific, Singapore, 2009).
- [13] J. Mathews and R. L. Walker, *Mathematical Methods of Physics* (W. A. Benjamin, Reading, MA, 1970).
- [14] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Perseus Books, Cambridge, Massachusetts, 1995).
- [15] C. Schubert, “Lectures on the worldline formalism”, in *School of Spinning Particles in Quantum Field Theory: Worldline Formalism, Higher Spins and Conformal Geometry*, edited by C. Schubert (2012).
- [16] M. Stone and P. Goldbart, *Mathematics for Physics: A Guided Tour for Graduate Students* (Cambridge Univ. Press, Cambridge, 2009).

Chapter 3

Daily QFT blog

2010-03-04 Predrag Kevin Mitchell is here, says we should study Littlejohn and M. Reinsch [39]: “Gauge fields in the separation of rotations and internal motions in the n -body problem.” I will put it into ChaosBook.org/library. Read 3-body problem sections. See p. 14 for a discussion of the three-body coordinates and p. 25 for a discussion of the three-body section (gauge).

2011-07-27 PC What follows is casting eye far ahead - to the role of gauge invariance in Quantum Field Theories. Following articles seem of interest as follow-ups on Cvitanović [11], *Group theory for Feynman diagrams in non-Abelian gauge theories*:

Khellat [34] strikes me as dubious...

Martens [43] writes: “We calculate the two-loop matching corrections for the gauge couplings at the Grand Unification scale in a general framework that aims at making as few assumptions on the underlying Grand Unified Theory (GUT) as possible. In this paper we present an intermediate result that is general enough to be applied to the Georgi-Glashow SU(5) as a “toy model”. The numerical effects in this theory are found to be larger than the current experimental uncertainty on α_s . Furthermore, we give many technical details regarding renormalization procedure, tadpole terms, gauge fixing and the treatment of group theory factors, which is useful preparative work for the extension of the calculation to supersymmetric GUTs.”

Tye and Zhang [61] write: “Bern, Carrasco and Johansson have conjectured dual identities inside the gluon tree scattering amplitudes. We use the properties of the heterotic string and open string tree scattering amplitudes to refine and derive these dual identities. These identities can be carried over to loop amplitudes using the unitarity method. Furthermore, given the M -gluon (as well as gluon-gluino) tree amplitudes, M -graviton (as well as graviton-gravitino) tree scattering amplitudes can be written down immediately, avoiding the derivation of Feynman rules and the evaluation of Feynman diagrams for graviton scattering amplitudes

Eto *et al.* [23] write: “We construct the general vortex solution in the color-flavor-locked vacuum of a non-Abelian gauge theory, where the gauge group is taken to be the product of an arbitrary simple group and $U(1)$. Use of the holomorphic invariants allows us to extend the moduli-matrix method and to determine the vortex moduli space in all cases. Our approach provides a new framework for studying solitons of non-Abelian varieties with various possible applications in physics.”

and there is much much more...; will continue some other time.

2011-11-03 PC Today is that time. I’m sitting in **Intractability Workshop: Counting, Inference and Optimization on Graphs** with a bunch of high-level computer nerds, and I almost afraid to say what I’ll say next (plumbers avoid physicists that say such things): In constructing our atlas of inertial manifold of turbulent pipe flow, we fix the $SO(2) \times O(2)$ phase separately on each local chart. The freedom of doing that is called “local gauge invariance” (blame Hermann Weyl for the ugly word) and in the limit of ∞ period cycles, cycle points are dense and their local charts are infinitesimal, so this is really local gauge invariance. In the world of computer science they use this freedom profitably, to reduce the number of terms they use in their computations. That suggests that there might be a (variational?) principle that selects an optimal choice of (relative) template phases (i.e., gauge transformations that connect a chart to the next chart).

Nerds call this ‘reparametrization’ - it supposedly speeds up calculations. Have not really seen that in quantum field theory, with exception of light cone gauges and their relatively recent applications by the Witten cult.

Literature: ref. [7, 8] and stuff on [this site](#) (if you can understand any of it).

Feel free to ignore this remark. It’s future research.

2012-05-16 Parameswaran Nair vpnair@optonline.net writes on saddle solutions of Yang-Mills:

I attributed the conjecture to Hitchin; it was actually due to Atiyah and Jones. “The only finite action solutions of the YM equations are instantons, either self-dual or anti-self-dual.” This was the conjecture for which the refs provide counter examples.

Here is the paper by my student Schiff [55], who writes: “ Following a proposal of Burzlaff (Phys.Rev.D 24 (1981) 546), we find solutions of the classical equations of motion of an abelian Higgs model on hyperbolic space, and thereby obtain a series of non-self-dual classical solutions of four-dimensional $SU(3)$ gauge theory. The lowest value of the action for these solutions is roughly 3.3 times the standard instanton action. ”

“ In physics, despite the fact that the non-self-dual solutions correspond to saddle points, and not minima, of the Yang-Mills functional, to do a correct semi-classical approximation by a saddle-point evaluation of the path integral, it is certainly necessary to include a contribution due to nonself-dual solutions, and if it should be the case that there is a non-self-dual solution with action lower

than the instanton action (this question is currently open, and of substantial importance), then such a contribution would even dominate. Unfortunately, it is questionable whether the semiclassical approximation can give a reliable picture of quantized gauge theories; it has been argued that in four-dimensional gauge theory small quantum fluctuations around classical solutions cannot be responsible for confinement, unlike in certain lower-dimensional theories. But it may still be possible to extract some physics from the semiclassical approach. A first step in such a direction would be to obtain a good understanding of the full set of non-self-dual solutions and their properties. [...] We pursue an old idea, due to Burzlaft [10], for obtaining a non-self-dual, "cylindrically symmetric" solution for gauge group $SU(3)$. If we write $\mathbb{R}^4 = \mathbb{R} \times \mathbb{R}^3$, and identify some $SU(2)$ [or $SO(3)$] subgroup of $SU(3)$, with generators that we will denote T , then we can look at the set of $SU(3)$ gauge potentials which are invariant under the action of the group generated by the sum of the T 's and the generators of rotations on the IR factor of E (we choose the T 's and the IR rotation generators to satisfy the same commutation relations). We call such potentials "cylindrically symmetric" (in analogy to the standard notion of cylindrical symmetry in IR, which involves writing $R = RXIR$ and requiring rotational symmetry on the E factor). Such potentials will be specified by a number of functions of two variables: the coordinate on the IR factor of E (which we will denote x), and the radial coordinate of the E factor (which we will denote y). Clearly the equations of motion for such cylindrically symmetric potentials (if they are consistent) will reduce to equations on the space $I(x,y); y \geq 0$.

The earlier work is by Sibner, Sibner, Uhlenbeck [58]. They write "The Yang-Mills functional for connections on principle $SU(2)$ bundles over S^4 is studied. Critical points of the functional satisfy a system of second-order partial differential equations, the Yang-Mills equations. If, in particular, the critical point is a minimum, it satisfies a first-order system, the self-dual or anti-self-dual equations. Here, we exhibit an infinite number of finite-action non-minimal unstable critical points. They are obtained by constructing a topologically nontrivial loop of connections to which min-max theory is applied. The construction exploits the fundamental relationship between certain invariant instantons on S^4 and magnetic monopoles on H^3 . This result settles a question in gauge field theory that has been open for many years."

Bor [2] writes "We prove the existence of a new family of non-self-dual finite-energy solutions to the Yang-Mills equations on Euclidean four-space, with $SU(2)$ as a gauge group. The approach is that of "equivariant geometry:" attention is restricted to a special class of fields, those that satisfy a certain kind of rotational symmetry, for which it is proved that (1) a solution to the Yang-Mills equations exists among them; and (2) no solution to the self-duality equations exists among them. The first assertion is proved by an application of the direct method of the calculus of variations (existence and regularity of minimizers), and the second assertion by studying the symmetry properties of the linearized self-duality equations. The same technique yields a new family of non-self-dual solutions on the complex projective plane."

2012-04-19 Daniel “graveyard of obvious ideas” rings a little aggressive, no? “if you are a master of quantum-mechanics or QFT symmetries and their linear irreducible representations, [14] you may leave your baggage at the door” rings a little aggressive, too.

Predrag: Get’s edgy. In “master of their linear irreducible representations” I make fun of myself. Let the referee object to that?

[2012-06-14 Predrag] Grin and bear it. Pulling my Senior discount card here.

2013-07-15 Predrag I’ve collected a bunch of QFT e-books, saved them in Chaos-Book.org/library:

5449Grigorenko06.pdf

Abarbanel 2013.pdf

CoKaWa04.pdf

DasFerb03.pdf

Zee03.pdf

hao89.pdf

Nichkawde13.pdf

Das06.pdf

Milton01.pdf

SeoSan12.pdf

Nagashima110.pdf

Elbaz12.pdf [19]

Sadovskii [54] *Quantum Field Theory*

Sadovskii [53] *Diagrammatics: Lectures on Selected Problems in Condensed Matter Theory*

2013-01-20 Predrag This really belongs to planar field theory, but for time being I note it here: Lucini and Panero [40] (in Chaosbook.org/library) might be of interest. All I get is one sentence and a reference only to ref. [13].

I should also read Kang and Loeb [33] *The enumeration of planar graphs via Wick’s theorem*.

2013-03-27 Predrag Do not understand this article: Jiménez-Lara and J. Llibre [29], *Periodic orbits and nonintegrability of generalized classical Yang–Mills Hamiltonian systems*.

Hu [28] *General initial value form of the semiclassical propagator*, write: “ We show a general initial value form of the semiclassical propagator. Similar to cellular dynamics, this formulation involves only the nearby orbits approximation: the evolution of nearby orbits is approximated by linearized dynamics. This phase space smearing formulation keeps the accuracy of the original Van Vleck-Gutzwiller propagator. As an illustration, we present a simple initial value form

of the semiclassical propagator. It is nonsingular everywhere and is efficient for numeric implementation. ”

2013-04-16 Predrag There seems to be whole literature on classical Yang-Mills (CYM). In *Entropy production in classical Yang-Mills theory from Glasma initial conditions* Hideaki Iida, Teiji Kunihiro, Berndt Müller, Akira Ohnishi, Andreas Schäfer, and Toru T. Takahashi, [arXiv:1304.1807](https://arxiv.org/abs/1304.1807), write:

Pure Yang-Mills theory in temporal gauge with the Hamiltonian in the noncompact (A, E) scheme on a cubic spatial lattice. The initial condition satisfies Gauss’ law; check its validity as well as Energy conservation carefully at every time step. Define distance (6), (7) that is gauge invariant under residual (time independent) gauge transformations.

They call the stability matrix ‘Hessian’, and its eigenvalues at time slice the ‘local Lyapunov exponents (LLEs)’ [38]: LLE plays the role of a “temporally local” Lyapunov exponent, which specifies the departure rate of two trajectories in a short time period. Then they say this (?): “For a system where stable and unstable modes couple with each other as in the present case, an LLE does not generally agree with the Lyapunov exponent in a long time period.” “Ref. [38] introduced another kind of Lyapunov exponent called the intermediate Lyapunov exponent (ILE), which is an “averaged Lyapunov exponent” for an intermediate time period; i.e., a time period which is sufficiently small compared to the thermalization time but large enough to sample a significant fraction of phase space. By its definition (13) it is the set of stability exponents for a finite time Jacobian matrix.

“Two comments are in order, here: A Lyapunov exponent [PC: not the Lyapunov exponent, they mean the stability exponent] can be (real) positive, negative, zero or purely imaginary. Liouville’s theorem tells us that the determinant of the time evolution matrix U is unity, implying that the sum of all positive and negative ILEs is zero. The KS entropy is given as a sum of positive Lyapunov exponents. The second comment concerns gauge invariance of the Lyapunov exponents. In the Appendix we show that LLE and ILE are indeed gauge invariant under time-independent gauge transformations in the temporal gauge.”

2013-11-27 Predrag One way to reduce symmetry seems obvious; **average over the group orbit**. That reduces the dimension of the state space by the dimension of the group orbit; in the reduced state space each group orbit is replaced by a point, it’s average value. It is a natural construct in the theory of linear representations of groups, very important, where it is called a ‘character’ of the representation; I use it in my [trace formula](#) for systems with continuous symmetries. But it is a trace of a linear evolution operator.

Still, I do not seem to know how to do this for (1) nonlinear systems, (2) QFT gauge fixing. Barth and Christensen [1] is an example where this is done - perhaps a way too complicated example... (here are my notes [on it](#), which I no longer understand myself :)

now in CB

2014-01-11 Predrag I always wonder whether we should be reducing symmetries by averaging over group orbits (method of characters, used in my derivation of the spectral determinant in presence of continuous symmetries). Churchill, Kummer and Rod [9] write in *On averaging, reduction, and symmetry in Hamiltonian systems*: The existence of periodic orbits for Hamiltonian systems at low positive energies can be deduced from the existence of nondegenerate critical points of an averaged Hamiltonian on an associated “reduced space.” The paper exploits discrete symmetries, including reversing diffeomorphisms, that occur in a given system. The symmetries are used to locate the periodic orbits in the averaged Hamiltonian, and thence in the original Hamiltonian when the periodic orbits are continued under perturbations admitting the same symmetries.”

2014-01-11 Predrag Kummer [37], *On the construction of the reduced phase space of a Hamiltonian system with symmetry* writes: “Weinstein [...] uses this correspondence between connections and lifts in his construction of Sternberg’s phase space for a particle in a Yang-Mills field. ”

1. A. Weinstein, A universal phase space for particles in Yang-Mills fields, *Lett. Math. Phys.* 2 (1978), 417-420.
2. S. STERNBERG, Minimal coupling and the symplectic mechanics of a classical particle in the presence of a Yang-Mills field, *Proc. Nat. Acad. Sci.* 74 (1977), 5253-5254.
3. S. STERNBERG, On the role of field theories in our physical conception of geometry in *Differential geometric methods in mathematical physics II*, Springer Lecture Notes in Mathematics, 676 (1977), 1-80.

2014-07-18 Predrag Mariño [42] lectures on non-perturbative effects are perhaps of interest: “a review of non-perturbative instanton effects in quantum theories, with a focus on large N gauge theories and matrix models. I first consider the structure of these effects in the case of ordinary differential equations, which provide a model for more complicated theories, and I introduce in a pedagogical way some technology from resurgent analysis, like trans-series and the resurgent version of the Stokes phenomenon. After reviewing instanton effects in quantum mechanics and quantum field theory, I address general aspects of large N instantons, and then present a detailed review of non-perturbative effects in matrix models.”

2014-07-30 Predrag Paul Hoyer recent talk at [Light Cone 2014](#) is too packed to be useful, but maybe has some pointers to recent interesting non-perturbative QCD results.

2014-08-01 Burak Sydney Coleman [10] gave a lecture on semi-classical formulations of QFT in 1975. He has several semi-classical treatment methods with names “Zeroth-Order Weak-Coupling Expansion”, “Coherent-State Variation”, “First-Order Weak-Coupling Expansion”, “Bohr-Sommerfeld Quantization”, and “DHN Formula”. I didn’t yet understand what any of these means, I’m just start-

ing to read it, but if we can find a way of adapting these methods to the numerical calculations that might be a good starting point.

2014-08-01 Predrag Coleman was a great expositor - on a level of a Nobelist. Would be cute if we found something in his lectures we can use today. "DHN Formula" (I knew all three - the first two authors are currently dead) should be the Gutzwiller formula for QFT. Paolo Muratore-Ginanneschi, a former student in our Copenhagen group, wrote something on that [48] that I still have not studied in depth. But we maybe should.

2014-08-01 Burak Coleman refers to the pedagogical overview by Rajaraman [51] for the derivation of "DHN Formula" and I think this will be a good starting point for me. So far, keywords are familiar, in the introduction, he talks about periodic orbits and their stability and how hard it is to find them for realistic cases. A lot of work apparently in this period has been done using Sine-Gordon equation because they had analytical solutions. I looked for articles which cite [51] but didn't see anything that relies on numerical solutions of field equations.

2014-08-13 Predrag In my notes it says that Dyson himself told me to read *Drawing theories apart: the dispersion of Feynman diagrams in postwar physics*, by Kaiser [31]. So we better read it - I will put a pilfered eBook copy [Kaiser09.pdf](#) into [ChaosBook.org/library](#).

2014-11-05 Predrag Do not know if this is something for us, but worth having a look at:

Luca Salasnich, *Discrete bright solitons in Bose-Einstein condensates and dimensional reduction in quantum field theory*, [arXiv:1411.0160](#): "We review the derivation of an effective one-dimensional (1D) discrete nonpolynomial Schrödinger equation from the continuous 3D Gross-Pitaevskii equation with transverse harmonic confinement and axial periodic potential. Then we study the bright solitons obtained from this discrete nonpolynomial equation showing that they give rise to the collapse of the condensate above a critical attractive strength. We also investigate the dimensional reduction of a bosonic quantum field theory, deriving an effective 1D nonpolynomial Heisenberg equation from the 3D Heisenberg equation of the continuous bosonic field operator under the action of transverse harmonic confinement. Moreover, by taking into account the presence of an axial periodic potential we find a generalized Bose-Hubbard model which reduces to the familiar 1D Bose-Hubbard Hamiltonian only if a strong inequality is satisfied. Remarkably, in the absence of axial periodic potential our 1D nonpolynomial Heisenberg equation gives the generalized Lieb-Liniger theory we obtained some years ago."

2014-11-23 Predrag Zvonkin [65] ([click here](#)) writes in *Matrix integrals and map enumeration: An accessible introduction*: "Physicists working in two-dimensional quantum gravity invented a new method of map enumeration based on computation of Gaussian integrals over the space of Hermitian matrices. This paper explains the basic facts of the method and provides an accessible introduction to the subject."

2015-01-09 Predrag Tudor Dimofte gave a talk in Math on *Geometric representation theory, symplectic duality, and 3d supersymmetric gauge theory*

Abstract: Recently, a “symplectic duality” between D-modules on certain pairs of algebraic symplectic manifolds was discovered, generalizing classic work of Beilinson-Ginzburg-Soergel in geometric representation theory. I will discuss how such dual spaces (some known and some new) arise naturally in supersymmetric gauge theory in three dimensions.

Tudor is a mathematical physicist at the IAS, School of Physics, Princeton.

I went to the talk, and - wow! You would think I know something about a gauge theory but it is like it was in Lithuanian: I understood individual words, and the alphabet seemed to be Latin - there were things that looked like letter G or letter H and what we call quotient M/G is apparently called ‘resolution’. The foundational paper is Braden, Licata, Proudfoot and Webster [5], and its followups on “Quantizations of conical symplectic resolutions II: category \mathcal{O} and symplectic duality”. Good luck reading these...

and of course, it was emphatically $N=4$ and not $N=2$, so now I’m at peace

:)

2015-02-04 Predrag Stephan Stetina <stetina@hep.itp.tuwien.ac.at> thesis on [arXiv:1502.00122](https://arxiv.org/abs/1502.00122) uses my [Field Theory](#), and says that 2PI graphs are no sweat (for me they were). Wrote to him:

You seem to have [proven Feynman wrong](#) :) That’s no mean achievement. Congrats!

Me and my friends have been studying turbulence in fluid dynamics as a warmup for doing the same in Yang-Mills. If you see some interesting turbulence in relativistic fluid dynamics, we are always willing to have a look at things more field-theoretical.

2015-02-09 Stephan Stetina If you are referring to Feynman comments on your book, I definitely disagree with them - I found your book on field theory more than helpful!

It is very difficult to study (quantum) turbulence within our approach - however it would be very interesting to do so! The original idea was to derive the two-fluid hydrodynamics of superfluids from an underlying field theory. To be able to obtain analytical results, we had to apply some rather drastic simplifications:

We assumed the superfluid condensate to be uniform and homogeneous (which translate in a homogeneous superflow in the hydro picture). Further more we used imaginary time formalism which strictly limits us to study systems in equilibrium. It would most likely be very challenging (in particular numerically) to introduce a condensate with arbitrary space and time dependence. In the current calculations, a probable onset of turbulence manifests itself as "something going wrong" - for example above certain velocities of the superfluid it is no longer possible to construct a stable and homogeneous superfluid phase. Another example is the appearance of the “two-stream instability” which can also be detected

in our approach (see for instance [arXiv:1312.5993](#)). I am not sure yet how much this approach has in common with the one you have cited.

2015-08-20 Predrag Strauss, Horwitz, Levitan, and Yahalom [60] *Quantum field theory of classically unstable Hamiltonian dynamics* might be a good starting point to learn about dynamical systems for which the motions can be described in terms of geodesics on a manifold. They say : “ ... ordinary potential models can be cast into this form by means of a conformal map. The geodesic deviation equation of Jacobi, constructed with a second covariant derivative, is unitarily equivalent to that of a parametric harmonic oscillator, and we study the second quantization of this oscillator. The excitations of the Fock space modes correspond to the emission and absorption of quanta into the dynamical medium, thus associating unstable behavior of the dynamical system with calculable fluctuations in an ensemble with possible thermodynamic consequences. ”

2016-01-08 Predrag Bogomolny wants us to study Englert and Schwinger [20–22]. Why? Ref. [22] *Semiclassical atom* seems to be reinventing Gutzwiller, without citing him. These papers are not cited much either, a pair of Nobel prizes notwithstanding :) Rohwedder and Englert [52] continue with *Semiclassical quantization in momentum space*. There is something called the Englert-Schwinger equation used in graphite studies [41]. Ullmo *et al.* [62] cite it: *Semiclassical density functional theory: Strutinsky energy corrections in quantum dots*

2016-04-06 Predrag Went to hear [Sung-Jin Oh](#) talk about

“... wave equations that arise from geometric considerations. Prime examples include the wave map equation and the Yang-Mills equation on the Minkowski space. On one hand, these are fundamental field theories arising in physics; on the other hand, they may be thought of as the hyperbolic analogues of the harmonic map and the elliptic Yang-Mills equations, which are interesting geometric PDEs on their own. I will discuss the recent progress on the problem of global regularity and asymptotic behavior of solutions to these PDEs.”

This kind of work might offer a path to computing non-trivial “exact coherent states” (non-perturbative classical solutions) of Yang-Mills. Experimentally I used Evernote on my phone to take notes and photos of the white board - [160406SungJinOh.pdf](#) in this repository - maybe it helps if one wants to get started reading the literature, though it is going to be hard going.

Both in the Abelian case (what they call “Maxwell-Klein-Gordon” for a scalar charged particle and “Maxwell-Dirac” for spin 1/2), and in the non-Abelian case (what they call “Yang-Mills”) they cheerfully set the particle mass to $m = 0$, which is a killer for us.

I tried to briefly explain to Sung-Jin Oh two things of possible interest to people solving the classical Maxwell-Klein-Gordon and Yang-Mills PDEs:

1. **finiteness conjecture**: perhaps they can find saddle-points (non-perturbative classical solutions of QED) that give the gauge-invariant sets as the starting step in perturbative calculations, rather than computing Feynman-diagrammatic corrections to trivial vacuum.

My explanation probably did more harm than good, as I liberally mixed in QFT, and kept confusing the quantum and the classical problem - it is for a reason that PDE people do not know what expressions like "on-mass-shell amplitudes" and "Ward identities" mean.

2. I ran Sung-Jin Oh quickly through ChaosBook.org/tutorials to make him aware that for turbulent nonlinear PDEs one has to work in the ∞ -dimensional state space, rather than look at the solutions only in the $(d+1)$ -dimensional configuration space.

He would have been more impressed if we could find such solutions for Eulerian flows, and give him a criterion which solutions are important, but I have no idea how to find smooth solutions for Euler (no viscosity Laplacian to help us there...).

All in all, I still have no idea for what kind of 'exact coherent states' to compute for Yang-Mills.

2016-05-26 Predrag William Graham Hoover and Kenichiro Aoki *Order and Chaos in the One-Dimensional ϕ^4 Model : N-Dependence and the Second Law of Thermodynamics*, [arXiv:1605.07721](https://arxiv.org/abs/1605.07721) write: " We revisit the equilibrium one-dimensional ϕ^4 model from the dynamical systems point of view. We find an infinite number of periodic orbits which are computationally stable while at the same time exhibiting positive Lyapunov exponents. We formulate a standard initial condition for the investigation of the microcanonical chaotic number dependence of the model. We speculate on the uniqueness of the model's chaotic sea and on the connection of such collections of deterministic and time-reversible states to the Second Law of Thermodynamics.

2016-10-12 Predrag Read Nguyen [49] *The perturbative approach to path integrals: A succinct mathematical treatment*: " We study finite-dimensional integrals in a way that elucidates the mathematical meaning behind the formal manipulations of path integrals occurring in quantum field theory. This involves a proper understanding of how Wick's theorem allows one to evaluate integrals perturbatively, i.e., as a series expansion in a formal parameter irrespective of convergence properties. We establish invariance properties of such a Wick expansion under coordinate changes and the action of a Lie group of symmetries, and we use this to study essential features of path integral manipulations, including coordinate changes, Ward identities, Schwinger-Dyson equations, Faddeev-Popov gauge-fixing, and eliminating fields by their equation of motion. We also discuss the asymptotic nature of the Wick expansion and the implications this has for defining path integrals perturbatively and nonperturbatively. "

2016-10-28 Predrag Read Hegg and Phillips [26] *Strongly coupled fixed point in φ^4 theory*: " We show explicitly how a fixed point can be constructed in scalar

$g\varphi^4$ theory from the solutions to a nonlinear eigenvalue problem. The fixed point is unstable and characterized by $\nu = 2/d$ (correlation length exponent), $\eta = 1/2 - d/8$ (anomalous dimension). For $d = 2$, these exponents reproduce to those of the Ising model which can be understood from the codimension of the critical point. The testable prediction of this fixed point is that the specific heat exponent vanishes. 2d critical Mott systems are well described by this new fixed point. ”

2016-12-10 Predrag Possibly useful in a QFT course:

Weinzierl [63] *Tales of 1001 Gluons*.

2015-09-15, 2017-02-13 Predrag Dashen, Hasslacher and Neveu [16–18], *Nonperturbative methods and extended-hadron models in field theory. I. Semiclassical functional methods*, are reputed to be the first people to use WKB methods in field theory.

Juan-Diego Urbina is a big fan of the 3rd paper [18]: “ a more modest approach by finding classical solutions of finite energy and bounded spatial extent. [...] We exhibit a four-dimensional model involving non-Abelian Yang-Mills fields. ”

Juan-Diego:

Ablowitz, Faddeev and Korepin have solitons for nonlinear Schrodinger on discrete lattice, with quartic term written as $|\psi_i|^2 \frac{1}{2}(\psi_{i-1} + \psi_{i+1})$. Even time can be taken discrete. Nohl took a one soliton solution, treated as a periodic orbit, got exact energy.

Soliton is an integrable, 4-parameter 4-dimensional manifold of solutions, in the infinite-dimensional space. All other action-angle pairs equal zero.

2016-12-26 Predrag Read Borinsky [3] *Renormalized asymptotic enumeration of Feynman diagrams*. It is a follow-up to Cvitanović, Lautrup and Pearson [15].

and Borinsky [4] PhD Thesis *Graphs in perturbation theory: Algebraic structure and asymptotics*

2017-05-26 Predrag Martin and Kearney [44] *An exactly solvable self-convolutive recurrence* fancy math reproduces (among much else) also counting of Cvitanović, Lautrup and Pearson [15] and Cvitanović [12]. The study the sequences $S(\alpha_1, \alpha_2, \alpha_3)$ of self-convolutive recurrences, derive a closed-form solutions as a Mellin transforms. The representation is useful for study the asymptotics via Laplace’s method. Their counting problem is the number of connected, or indecomposable, permutations, which naturally leads them to QFT diagram counting; For example, the number of nonisomorphic connected Feynman diagrams of order $2(n + 1)$ arising in a simplified model of quantum electrodynamics (QED) [15]; the number of ‘vertex graphs’ of order $2n$ arising in the QED perturbation series for the electron magnetic moment [12]; the number of Feynman diagrams with exact propagators [12]; and the number of Feynman diagrams with proper self-energies arising in QED [15]. Their integral representation is apparently new.

Smith [59] *Planar version of Baym-Kadanoff theory*

Kugler [36] *Counting Feynman diagrams via many-body relations*

Kaneko [32] *Enumeration of N-rooted maps using quantum field theory*: “ Information about the number of Feynman graphs for a given physical process in a given field theory is especially useful for confirming the result of a Feynman graph generator used in an automatic system of perturbative calculations. A method of counting the number of Feynman graphs with weight of symmetry factor was established based on zero-dimensional field theory, and was used in scalar theories and QED. In this article this method is generalized to more complicated models by direct calculation of generating functions on a symbolic calculating system. This method is applied to QCD with and without counter terms, where many higher order are being calculated automatically. ”

2017-05-29 Predrag Read Pavlyukh and W. Hübner [50] *Analytic solution of Hedin’s equations in zero dimensions*: “ Feynman diagrams for the many-body perturbational theory are enumerated by solving the system of Hedin’s equations in zero dimension. We extend the treatment of Molinari [45] and give a complete solution of the enumeration problem in terms of Whittaker functions. An important relation between the generating function of the electron propagator and anomalous dimension in quantum field theory of massless fermions and mesons in four dimensions (Yukawa theory) is found. The Hopf algebra of undecorated rooted trees yields the anomalous field dimension in terms of the solution of the same differential equation. Its relation to the mathematical problem of combinatorics of chord diagrams is discussed; asymptotic expansions of the counting numbers are obtained. ”

2018-04-28 Predrag Castro and Roditi [6] *A combinatorial matrix approach for the generation of vacuum Feynman graphs multiplicities in ϕ^4 theory*. They cite Cvitanović, Lautrup and Pearson [15] and write: “ [...] generate the set of all Feynman graphs and the respective multiplicities in a combinatoric way. These combinatorial matrices are explicitly related with the permutation group, which facilitates the construction of the vacuum Feynman graphs. Various insights in this combinatoric problem are proposed, which in principle provide an efficient way to compute the Feynman vacuum graphs and its multiplicities. ”

Gopala, Labelle and Shramchenko [30] *Enumeration of N-rooted maps using quantum field theory* prove, inter alia, the equality of the number of two-point Feynman diagrams in scalar QED [15] and the number of rooted maps.

2017-05-29 Predrag Read Molinari [45, 46] *Hedin’s equations and enumeration of Feynman diagrams*

Molinari and N. Manini [47] *Enumeration of many-body skeleton diagrams* “Based on Hedin’s equations for self-energy, polarization, propagator, effective potential, and vertex function, dressed (skeleton) Feynman diagrams are enumerated.”

2017-05-29 Predrag Read Krishna Gopala, Labelle and Shramchenko [35] *Enumeration of N-rooted maps using quantum field theory*: “ A one-to-one correspondence is proved between the N-rooted ribbon graphs, or maps, with e edges and

the $(e-N+1)$ -loop Feynman diagrams of a certain quantum field theory. This result is used to obtain explicit expressions and relations for the generating functions of N -rooted maps and for the numbers of N -rooted maps with a given number of edges using the path integral approach. ”

2017-03-15 Predrag Read Setlur [57] *Dynamics of Classical and Quantum Fields: An Introduction*. Starts with Geometrical meaning of Legendre transformation in classical mechanics; Dynamical symmetries in the context of Noether’s theorem; The derivation of the stress energy tensor of the electromagnetic field; the expression for strain energy in elastic bodies, and the Navier Stokes equation; Functional integration is interpreted as a limit of a sequence of ordinary integrations, The rest is less obvious.

In principle, this book can be read online via `library.gatech.edu`, but how?

2018-01-31 Predrag Check out M. J. D. Hamilton [25] *Mathematical Gauge Theory*: “ This book explains the mathematical background behind the Standard Model, translating ideas from physics into a mathematical language and vice versa. The first part of the book covers the mathematical theory of Lie groups and Lie algebras, fibre bundles, connections, curvature and spinors. The second part then gives a detailed exposition of how these concepts are applied in physics, concerning topics such as the Lagrangians of gauge and matter fields, spontaneous symmetry breaking, the Higgs boson and mass generation of gauge bosons and fermions. [...] Only a basic knowledge of differentiable manifolds and special relativity is required, summarized in the appendix. ”

2018-03-31 Predrag Created the abstract for the Les Houches meeting, see `predrag/lectures/LesHouch18`. The talk is in this repo, `reducesymm/presentations/LesHouch18`
The worldline formula for the quenched QED form factors is currently wrong; essentially it puts Z_2 renormalization on each electron leg, rather than $\sqrt{Z_2}$.

2018-06-05 Predrag Notes on *Summer school on structures in local quantum field theory Les Houches* — June 4-15, 2018

Gerald Dunne *Resurgence and Perturbative/Non-Perturbative Relations* is pretty amazing - exact expression for trace formula in terms of WKB saddles and the non-perturbative corrections, based on 2-torus duality relations, see [here](#) for lecture notes (I have also written down some notes). I think this works only for integrable models. Told him to have a look at Wirzba [64] as a physically motivated chaotic problem whose analytic formulation is suited to Dunne’s methods.

Jacob Bourjaily *Improving Integrands and Integrals for Amplitudes*: It’s complicated. However, I did explain to him how to reduce all adjoint rep birdtracks (quotient Jacobi relation, treat the rest as a free algebra) to a basis set of trees + fully fully symmetric Casimir tensors; he should get back to me once he tries it.

2018-06-06 Predrag **James M. Drummond** *Cluster algebras and scattering amplitudes*, [arXiv:1710.10953](#), was a stunning performance, a picture of which I’ll

post on Flickr.com. Analytic relations for scattering amplitudes expressed the cluster algebras for planar $N = 4$ super Yang-Mills theory that generalize the Steinmann relations. I still do not know what hit me.

2018-06-07 Predrag Lorenzo Magnea (an intellectual, with opinions on many things, masquerading as a phenomenologist, five layers removed from LHC experimentalists) *Eikonal Correlators and form factors in perturbation theory*, and **Franz Herzog** *Geometric IR subtraction in real radiation*, [arXiv:1804.07949](#), are our best shot for developing approximations to N -photon propagators to all orders. The idea is to claim that the N -photon propagator in the “rainbow” gauge sets a_{N00} is concentrated on (Schwinger) backbone that hops over the magnetic momentum vertex with a large momentum q , with all photons ‘soft’ respective to the backbone, with momenta $q - k_j$, $|k_j| \ll 1$. Then the same discussion as for IR contributions should apply, with only one γ^μ per N -photon propagator mass-shell vertex surviving in the IR limit, the rest reducing to scalar, gauge invariant and universal M_{soft} , commuting vertices, and exponentiating. The limit is explained in Magnea *Advanced Lectures on the Infrared structure of Perturbative QCD* handwritten [lecture 2](#), sect. 2 *The soft approximation*, and the all order summation is given in [lecture 3](#).

Study also Herzog [\[27\]](#) *Zimmermann’s forest formula, infrared divergences and the QCD beta function*.

The trick then would be to formulate next-to-leading order (NLO) and next-to-next-to-leading order (NNLO) corrections systematically.

Henry Kißler *The t’Hooft–Veltman gauge*: Finally understood the point of Henry’s thesis. The t’Hooft–Veltman gauge condition is quadratic in A^μ , leading to 4-vertices, and QED ghost sector with structure reminiscent of a non-Abelian QCD. By Ward identities it all adds up to zero, but individual graphs are non-trivial.

Erik Panzer *The Hepp bound for Feynman periods*. “Feynman period” is this cult’s jargon for the value of a Feynman integral, a quantity hard to evaluate except for the known zoo of integrals evaluated so far, catalogued by **Oliver Schnetz** [\[56\]](#). The “Hepp bound” or “Hepp invariant,” however, is a *rational* number computed efficiently for all graphs. Some of the striking properties of this bound: it correlates very well with the actual Feynman period (the empirical ratio of the two fluctuates within 1%, a numerical fact not explained yet) and it respects all known identities among periods (I have some handwritten notes on this talk).

From gauge-sets point of view, analytic formulas for (g-2) seem useless, as they are sets of large exact numbers (see eq. (29) in Schnetz [\[56\]](#); [arXiv:1711.05118](#)), that then sum up to a much smaller number.

I had extensive discussion with Erik, and gauge sets are now in his “to-do” list. While Schnetz and Panzer had focused on ϕ^4 , and I have this deep prejudice that gauge theories are profoundly different (because I assume that the gauge invariance induces cancelations among finite parts), perhaps they are not so different.

Let's think of gauge symmetry as we think of other continuous symmetries in theory of dynamical systems. Then quotienting the symmetry leads us to gauge sets, which have no symmetry, but each has $n!$ Feynman diagram contributions - a " ϕ^4 " for each gauge set. Erik tells me that someone has proven that Borel-resummed ϕ^4 has a finite radius of convergence, so it still could work out that the quenched QED has a finite radius of convergence.

2018-06-08 Predrag Spencer Bloch *The algebraic geometry of the kite graph amplitude* assumed more pre-knowledge than what I possess. Like I've spent last 20 years chatting to Spencer, and he's telling me what's new in the last 6 weeks, without defining a single thing.

Berghoff

2018-06-11 Predrag Dirk Kreimer *Complex graphs and graph complexes*

Schlotterer

Marcel Golz *Parametric QED* [24]. The Schwinger parametric Feynman integrals for gauge theories quickly become prohibitively complicated due to the very involved numerator polynomials. He reported on a simplification via combinatorics for QED; the integrand is only a very small sum of scalar integrands with Dodgson polynomial (introduced by Francis Brown) numerators. Dodgson polynomial (related to cycle polynomials) satisfy Dodgson identities. Marcel is interested in simplifications such as organization by gauge sets, so I gave him access to this blog.

Bogner

Brödel

Jaclyn Bell is a charismatic young Liverpoolsian (from working class background), theoretical particle physics PhD advised by a Northern Irish childhood friend and colleague of Georgia Tech's Brian Kennedy, who did a stint on the BBC astronauts show as potential astronaut, and now runs a UK STEM education organization, with outreach to children in UK's poorer neighborhoods. If I got it right....

2018-06-12 Predrag Wulkenhaar (Meintz) had a breakthrough assist from Erik Panzer during this workshop, which seems to establish that a non-commutative matrix ϕ^4 model is integrable. (My mother's ϕ^4 QFT model remains definitely non-integrable.) I have handwritten notes on his talk.

Alexander Hock *Noncommutative 3-colour scalar quantum field theory model in 2D* is a closely related model with the same structure: "We introduce the 3-colour noncommutative quantum

field theory model in two dimensions. For this model we prove a generalised Ward-Takahashi identity, which is special to coloured noncommutative QFT models. It reduces to the usual Ward-Takahashi identity in a particular case. The Ward-Takahashi identity is used to simplify the Schwinger-Dyson equations for the 2-point function and the N -point function. The absence of any renormalisation conditions in the large (N, V) -limit in 2D leads to a recursive integral

equation for the 2-point function, which we solve perturbatively to sixth order in the coupling constant. One important fact is the appearance of polylogarithms in the perturbative solution, which is generated by the closed integral equation. With the knowledge of the experts here in Les Houches I hope to improve my results to higher order or even

and an exact solution for the 2-point function. ”

Masha Vlasenko *Motivic Gamma Functions*, see “what is a [motivic gamma function?](#)”. Not a chance that I would understand anything.

2018-06-13 Predrag Burgos-Gil is a lovely Spanish-speaking refugee from Barcelona living in Madrid.

Ralph Kaufmann *Feynman categories and applications: geometry, number theory and physics*

David Prinz *Einstein-Maxwell-Dirac theory* is the canonical generalization of spinor electrodynamics to curved spacetimes of general relativity. He explains the underlying geometry of the theory and its Lagrange density, with gauge fixing, ghost terms, Feynman rules and tree-level interactions. The obstructions to multiplicative renormalization are overcome by a generalization of Furry’s theorem.

2018-06-16 Predrag Remember to send Henry Kibler a hard copy of my “Field Theory” book(lett).

References

- [1] N. H. Barth and S. M. Christensen, “[Quantizing fourth-order gravity theories: The functional integral](#)”, *Phys. Rev. D* **28**, 1876–1893 (1983).
- [2] G. Bor, “[Yang-Mills fields which are not self-dual](#)”, *Commun. Math. Phys.* **145**, 393–410 (1992).
- [3] M. Borinsky, “[Renormalized asymptotic enumeration of Feynman diagrams](#)”, *Annals of Physics* (2017) [10.1016/j.aop.2017.07.009](#).
- [4] M. Borinsky, [Graphs in perturbation theory: Algebraic structure and asymptotics](#), PhD thesis (Humboldt-Universität, Berlin, 2018).
- [5] T. Braden, A. Licata, N. Proudfoot, and B. Webster, “[Hypertoric category](#)”, *Adv. Math.* **231**, 1487–1545 (2012).
- [6] E. Castro and I. Roditi, [Enumeration of n-rooted maps using quantum field theory](#), 2018.
- [7] V. Y. Chernyak and M. Chertkov, “[Fermions and loops on graphs: I. Loop calculus for determinants](#)”, *J. Stat. Mech.* **2008**, P12011 (2008).
- [8] M. Chertkov and A. B. Yedidia, “[Approximating the permanent with fractional belief propagation](#)”, *J. Machine Learning Research* **14**, 2029–2066 (2013).

- [9] R. C. Churchill, M. Kummer, and D. L. Rod, “On averaging, reduction, and symmetry in Hamiltonian systems”, *J. Diff. Eqn.* **49**, 359–414 (1983).
- [10] S. Coleman, “Classical lumps and their quantum descendants”, in *New Phenomena in Subnuclear Physics*, Vol. 13, edited by A. Zichichi (Springer, 1977), pp. 297–421.
- [11] P. Cvitanović, “Group theory for Feynman diagrams in non-Abelian gauge theories”, *Phys. Rev. D* **14**, 1536–1553 (1976).
- [12] P. Cvitanović, “Asymptotic estimates and gauge invariance”, *Nucl. Phys. B* **127**, 176–188 (1977).
- [13] P. Cvitanović, “Planar perturbation expansion”, *Phys. Lett. B* **99**, 49–52 (1981).
- [14] P. Cvitanović, *Group Theory: Birdtracks, Lie’s and Exceptional Groups* (Princeton Univ. Press, Princeton NJ, 2004).
- [15] P. Cvitanović, B. Lautrup, and R. B. Pearson, “Number and weights of Feynman diagrams”, *Phys. Rev. D* **18**, 1939–1949 (1978).
- [16] R. F. Dashen, B. Hasslacher, and A. Neveu, “Nonperturbative methods and extended-hadron models in field theory. I. Semiclassical functional methods”, *Phys. Rev. D* **10**, 4114–4129 (1974).
- [17] R. F. Dashen, B. Hasslacher, and A. Neveu, “Nonperturbative methods and extended-hadron models in field theory. II. Two-dimensional models and extended hadrons”, *Phys. Rev. D* **10**, 4130–4138 (1974).
- [18] R. F. Dashen, B. Hasslacher, and A. Neveu, “Nonperturbative methods and extended-hadron models in field theory. III. Four-dimensional non-Abelian models”, *Phys. Rev. D* **10**, 4138–4142 (1974).
- [19] E. Elbaz, *Quantum: The Quantum Theory of Particles, Fields and Cosmology* (Springer, Berlin, 2012).
- [20] B.-G. Englert and J. Schwinger, “Atomic-binding-energy oscillations”, *Phys. Rev. A* **32**, 47–63 (1985).
- [21] B.-G. Englert and J. Schwinger, “Linear degeneracy in the semiclassical atom”, *Phys. Rev. A* **32**, 36–46 (1985).
- [22] B.-G. Englert and J. Schwinger, “Semiclassical atom”, *Phys. Rev. A* **32**, 26–35 (1985).
- [23] M. Eto, T. Fujimori, S. B. Gudnason, K. Konishi, M. Nitta, K. Ohashi, and W. Vinci, “Constructing non-Abelian vortices with arbitrary gauge groups”, *Phys. Lett. B* **669**, 98–101 (2008).
- [24] M. Golz, “New graph polynomials in parametric QED feynman integrals”, *Ann. Phys.* **385**, 328–346 (2017).
- [25] M. J. Hamilton, *Mathematical Gauge Theory* (Springer, New York, 2017).
- [26] A. Hegg and P. W. Phillips, “Strongly coupled fixed point in φ^4 theory”, *Europhys. Lett.* **115**, 27005 (2016).

- [27] F. Herzog, “Zimmermann’s forest formula, infrared divergences and the QCD beta function”, *Nucl. Phys. B* **926**, 370–380 (2018).
- [28] B. Hu, Q. Jie, B. Li, and S. Wang, “General initial value form of the semiclassical propagator”, *Phys. Rev. A* **63**, 044102 (2001).
- [29] L. Jiménez-Lara and J. Llibre, “Periodic orbits and nonintegrability of generalized classical Yang–Mills Hamiltonian systems”, *J. Math. Phys.* **52**, 032901 (2011).
- [30] V. S. K. K. Gopala P. Labelle, *Enumeration of N-rooted maps using quantum field theory*, 2017.
- [31] D. Kaiser, *Drawing Theories Apart: The Dispersion of Feynman Diagrams in Postwar Physics* (Univ. Chicago Press, 2009).
- [32] T. Kaneko, *Counting the number of Feynman graphs in QCD*, 2017.
- [33] M. Kang and M. Loeb, “The enumeration of planar graphs via Wick’s theorem”, *Adv. Math.* **221**, 1703–1724 (2009).
- [34] M. Khellat and A. Mirjalili, *Finding patterns within perturbative approximation in QCD and indirect relations*, 2010.
- [35] K. Krishna Gopala, P. Labelle, and V. Shramchenko, *Enumeration of N-rooted maps using quantum field theory*, 2017.
- [36] F. B. Kugler, *Counting Feynman diagrams via many-body relations*, 2018.
- [37] M. Kummer, “On the construction of the reduced phase space of a Hamiltonian system with symmetry”, *Indiana Univ. Math. J* **30**, 281–291 (1981).
- [38] T. Kunihiro, B. Müller, A. Ohnishi, A. Schäfer, T. T. Takahashi, and A. Yamamoto, “Chaotic behavior in classical Yang–Mills dynamics”, *Phys. Rev. D* **82**, 114015 (2010).
- [39] R. G. Littlejohn and M. Reinsch, “Gauge fields in the separation of rotations and internal motions in the n -body problem”, *Rev. Mod. Phys.* **69**, 213–275 (1997).
- [40] B. Lucini and M. Panero, “SU(N) gauge theories at large N”, *Phys. Rep.* **526**, 93–163 (2013).
- [41] A. Mañanes and E. Santos, “Interplanar binding in graphite studied with the Englert–Schwinger equation”, *Phys. Rev. B* **34**, 5874–5882 (1986).
- [42] M. Mariño, “Lectures on non-perturbative effects in large N gauge theories, matrix models and strings”, *Fortschr. Phys.* **62**, 455–540 (2014).
- [43] W. Martens, “Towards a two-loop matching of gauge couplings in grand unified theories”, *J. High Energy Phys.* **2011**, 1–35 (2011).
- [44] R. J. Martin and M. J. Kearney, “An exactly solvable self-convolutive recurrence”, *Aequat. Math.* **80**, 291–318 (2010).
- [45] L. G. Molinari, “Hedin’s equations and enumeration of Feynman diagrams”, *Phys. Rev. B* **71**, 113102 (2005).

- [46] L. G. Molinari, “A derivation of Hedin’s equations”, *Phys. Rev. B* **71**, 113102 (2007).
- [47] L. G. Molinari and N. Manini, “Enumeration of many-body skeleton diagrams”, *Eur. Phys. J. B* **51**, 331–336 (2006).
- [48] P. Muratore-Ginanneschi, “Path integration over closed loops and Gutzwiller’s trace formula”, *Phys. Rep.* **383**, arXiv:nlin/0210047, 299–397 (2002).
- [49] T. Nguyen, “The perturbative approach to path integrals: A succinct mathematical treatment”, *J. Math. Phys.* **57**, 092301 (2016).
- [50] Y. Pavlyukh and W. Hübner, “Analytic solution of Hedin’s equations in zero dimensions”, *J. Math. Phys.* **48**, 052109 (2007).
- [51] R. Rajaraman, “Some non-perturbative semi-classical methods in quantum field theory (a pedagogical review)”, *Physics Reports* **21**, 227–313 (1975).
- [52] B. Rohwedder and B.-G. Englert, “Semiclassical quantization in momentum space”, *Phys. Rev. A* **49**, 2340–2346 (1994).
- [53] M. V. Sadovskii, *Diagrammatics: Lectures on Selected Problems in Condensed Matter Theory* (World Scientific, Singapore, 2006).
- [54] M. V. Sadovskii, *Quantum Field Theory* (De Gruyter, 2013).
- [55] J. Schiff, “Hyperbolic vortices and some non-self-dual classical solutions of SU(3) gauge theory”, *Phys. Rev. D* **44**, 528–535 (1991).
- [56] O. Schnetz, “Numbers and functions in quantum field theory”, *Phys. Rev. D* **97**, 085018 (2018).
- [57] G. Setlur, *Dynamics of Classical and Quantum Fields: An Introduction* (CRC Press, Boca Raton, FL, 2013).
- [58] L. M. Sibner, R. J. Sibner, and K. Uhlenbeck, “Solutions to Yang-Mills equations that are not self-dual”, *Proc. Natl. Acad. Sci. USA* **86**, 8610–8613 (1989).
- [59] R. A. Smith, “Planar version of Baym-Kadanoff theory”, *Phys. Rev. A* **46**, 4586–4597 (1992).
- [60] Y. Strauss, L. P. Horwitz, J. Levitan, and A. Yahalom, “Quantum field theory of classically unstable Hamiltonian dynamics”, *J. Math. Phys.* **56**, 072701 (2015).
- [61] S.-H. H. Tye and Y. Zhang, “Dual identities inside the gluon and the graviton scattering amplitudes”, *J. High Energy Phys.* **2010**, 1–45 (2010).
- [62] D. Ullmo, T. Nagano, S. Tomsovic, and H. U. Baranger, “Semiclassical density functional theory: Strutinsky energy corrections in quantum dots”, *Phys. Rev. B* **63**, 125339 (2001).
- [63] S. Weinzierl, “Tales of 1001 Gluons”, *Physics Reports* **676**, 1–101 (2017).
- [64] A. Wirzba, “Quantum mechanics and semiclassics of hyperbolic n-disk scattering systems”, *Phys. Rep.* **309**, 1–116 (1999).
- [65] A. Zvonkin, “Matrix integrals and map enumeration: An accessible introduction”, *Math. Comput. Modelling* **26**, 281–304 (1997).